

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
25 April 2002 (25.04.2002)

PCT

(10) International Publication Number  
**WO 02/33060 A2**

(51) International Patent Classification<sup>7</sup>: C12N 9/00

(74) Agent: MARSH, David, R.; Arnold & Porter, 555 12th Street, N.W., Washington, DC 20004 (US).

(21) International Application Number: PCT/US01/42673

(22) International Filing Date: 12 October 2001 (12.10.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
09/688,071 14 October 2000 (14.10.2000) US

(71) Applicant: MONSANTO TECHNOLOGY LLC  
[US/US]; 800 N. Lindbergh Blvd., St. Louis, MO 63167 (US).

(72) Inventors: LASSNER, Michael, W.; 721 Falcon Avenue, Davis, CA 95616 (US). SAVIDGE, Beth; 3212 Chesapeake Bay Avenue, Davis, CA 95616 (US). WEISS, James, D.; 471 Goethe Avenue, Kirkwood, MO 63122 (US). MITSKY, Timothy, A.; 2262 A Rule Avenue, Maryland Heights, MO 63043 (US). POST-BEITTEN-MILLER, Martha, Ann; 601 Lalor Drive, Manchester, MO 63011 (US). VALENTIN, Henry, E.; 873 M Fox Spring Dr., Chesterfield, MO 63017 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

**Published:**

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

WO 02/33060 A2

(54) Title: NUCLEIC ACID SEQUENCES TO PROTEINS INVOLVED IN TOCOPHEROL SYNTHESIS

(57) Abstract: Nucleic acid sequences and methods are provided for producing plants and seeds having altered tocopherol content and compositions. The methods find particular use in increasing the tocopherol levels in plants, and in providing desirable tocopherol compositions in a host plant cell.

## NUCLEIC ACID SEQUENCES TO PROTEINS INVOLVED IN TOCOPHEROL SYNTHESIS

5

### INTRODUCTION

This application claims the benefit of the filing date of US. Application Serial Number 09/549,848, filed April 14, 2000.

10

### TECHNICAL FIELD

The present invention is directed to nucleic acid and amino acid sequences and constructs, and methods related thereto.

15

### BACKGROUND

Isoprenoids are ubiquitous compounds found in all living organisms. Plants synthesize a diverse array of greater than 22,000 isoprenoids (Connolly and Hill (1992) *Dictionary of Terpenoids*, Chapman and Hall, New York, NY). In plants, isoprenoids play essential roles in particular cell functions such as production of sterols, contributing to eukaryotic membrane architecture, acyclic polyprenoids found in the side chain of ubiquinone and plastoquinone, growth regulators like abscisic acid, gibberellins, brassinosteroids or the photosynthetic pigments chlorophylls and carotenoids. Although the physiological role of other plant isoprenoids is less evident, like that of the vast array of secondary metabolites, some are known to play key roles mediating the adaptative responses to different environmental challenges. In spite of the remarkable diversity of structure and function, all isoprenoids originate from a single metabolic precursor, isopentenyl diphosphate (IPP) (Wright, (1961) *Annu. Rev. Biochem.* 20:525-548; and Spurgeon and Porter, (1981) in Biosynthesis of Isoprenoid Compounds, Porter and Spurgeon eds (John Wiley, New York) Vol. 1, pp1-46).

20

25

A number of unique and interconnected biochemical pathways derived from the isoprenoid pathway leading to secondary metabolites, including tocopherols, exist in chloroplasts

30

of higher plants. Tocopherols not only perform vital functions in plants, but are also important from mammalian nutritional perspectives. In plastids, tocopherols account for up to 40% of the total quinone pool.

5 Tocopherols and tocotrienols (unsaturated tocopherol derivatives) are well known antioxidants, and play an important role in protecting cells from free radical damage, and in the prevention of many diseases, including cardiac disease, cancer, cataracts, retinopathy, Alzheimer's disease, and neurodegeneration, and have been shown to have beneficial effects on symptoms of arthritis, and in anti-aging. Vitamin E is used in chicken feed for improving the shelf life, appearance, flavor, and oxidative stability of meat, and to transfer tocopherols from feed to  
10 eggs. Vitamin E has been shown to be essential for normal reproduction, improves overall performance, and enhances immunocompetence in livestock animals. Vitamin E supplement in animal feed also imparts oxidative stability to milk products.

The demand for natural tocopherols as supplements has been steadily growing at a rate of 10-20% for the past three years. At present, the demand exceeds the supply for natural  
15 tocopherols, which are known to be more biopotent than racemic mixtures of synthetically produced tocopherols. Naturally occurring tocopherols are all *d*-stereoisomers, whereas synthetic  $\alpha$ -tocopherol is a mixture of eight *d,l*- $\alpha$ -tocopherol isomers, only one of which (12.5%) is identical to the natural *d*- $\alpha$ -tocopherol. Natural *d*- $\alpha$ -tocopherol has the highest vitamin E activity (1.49 IU/mg) when compared to other natural tocopherols or tocotrienols. The synthetic  $\alpha$ -tocopherol  
20 has a vitamin E activity of 1.1 IU/mg. In 1995, the worldwide market for raw refined tocopherols was \$1020 million; synthetic materials comprised 85-88% of the market, the remaining 12-15% being natural materials. The best sources of natural tocopherols and tocotrienols are vegetable oils and grain products. Currently, most of the natural Vitamin E is produced from  $\gamma$ -tocopherol derived from soy oil processing, which is subsequently converted to  
25  $\alpha$ -tocopherol by chemical modification ( $\alpha$ -tocopherol exhibits the greatest biological activity).

Methods of enhancing the levels of tocopherols and tocotrienols in plants, especially levels of the more desirable compounds that can be used directly, without chemical modification, would be useful to the art as such molecules exhibit better functionality and bioavailability.

In addition, methods for the increased production of other isoprenoid derived compounds in a host plant cell is desirable. Furthermore, methods for the production of particular isoprenoid compounds in a host plant cell is also needed.

5

## SUMMARY OF THE INVENTION

The present invention is directed to sequences to proteins involved in tocopherol synthesis. The polynucleotides and polypeptides of the present invention include those derived from prokaryotic and eukaryotic sources.

10

Thus, one aspect of the present invention relates to prenyltransferase, and in particular to isolated polynucleotide sequences encoding prenyltransferase proteins and polypeptides related thereto. In particular, isolated nucleic acid sequences encoding prenyltransferase proteins from bacterial and plant sources are provided.

15

In another aspect, the present invention provides isolated polynucleotide sequences encoding tocopherol cyclase, and polypeptides related thereto. In particular, isolated nucleic acid sequences encoding tocopherol cyclase proteins from bacterial and plant sources are provided.

Another aspect of the present invention relates to oligonucleotides which include partial or complete prenyltransferase or tocopherol cyclase encoding sequences.

20

It is also an aspect of the present invention to provide recombinant DNA constructs which can be used for transcription or transcription and translation (expression) of prenyltransferase or tocopherol cyclase. In particular, constructs are provided which are capable of transcription or transcription and translation in host cells.

25

In another aspect of the present invention, methods are provided for production of prenyltransferase or tocopherol cyclase in a host cell or progeny thereof. In particular, host cells are transformed or transfected with a DNA construct which can be used for transcription or transcription and translation of prenyltransferase or tocopherol cyclase. The recombinant cells which contain prenyltransferase or tocopherol cyclase are also part of the present invention.

30

In a further aspect, the present invention relates to methods of using polynucleotide and polypeptide sequences to modify the tocopherol content of host cells, particularly in host plant



cells. Plant cells having such a modified tocopherol content are also contemplated herein.

Methods and cells in which both prenyltransferase and tocopherol cyclase are expressed in a host cell are also part of the present invention.

The modified plants, seeds and oils obtained by the expression of the prenyltransferase or  
5 tocopherol cyclase are also considered part of the invention.

### BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 provides an amino acid sequence alignment between ATPT2, ATPT3, ATPT4,  
10 ATPT8, and ATPT12 are performed using ClustalW.

Figure 2 provides a schematic picture of the expression construct pCGN10800.

Figure 3 provides a schematic picture of the expression construct pCGN10801.

Figure 4 provides a schematic picture of the expression construct pCGN10803.

Figure 5 provides a schematic picture of the construct pCGN10806.

15 Figure 6 provides a schematic picture of the construct pCGN10807.

Figure 7 provides a schematic picture of the construct pCGN10808.

Figure 8 provides a schematic picture of the expression construct pCGN10809.

Figure 9 provides a schematic picture of the expression construct pCGN10810.

Figure 10 provides a schematic picture of the expression construct pCGN10811.

20 Figure 11 provides a schematic picture of the expression construct pCGN10812.

Figure 12 provides a schematic picture of the expression construct pCGN10813.

Figure 13 provides a schematic picture of the expression construct pCGN10814.

Figure 14 provides a schematic picture of the expression construct pCGN10815.

Figure 15 provides a schematic picture of the expression construct pCGN10816.

25 Figure 16 provides a schematic picture of the expression construct pCGN10817.

Figure 17 provides a schematic picture of the expression construct pCGN10819.

Figure 18 provides a schematic picture of the expression construct pCGN10824.

Figure 19 provides a schematic picture of the expression construct pCGN10825.

Figure 20 provides a schematic picture of the expression construct pCGN10826.

Figure 21 provides an amino acid sequence alignment using ClustalW between the *Synechocystis* prenyltransferase sequences.

Figure 22 provides an amino acid sequence of the ATPT2, ATPT3, ATPT4, ATPT8, and ATPT12 protein sequences from *Arabidopsis* and the slr1736, slr0926, slr1899, slr0056, and the  
5 slr1518 amino acid sequences from *Synechocystis*.

Figure 23 provides the results of the enzymatic assay from preparations of wild type *Synechocystis* strain 6803, and *Synechocystis* slr1736 knockout.

Figure 24 provides bar graphs of HPLC data obtained from seed extracts of transgenic *Arabidopsis* containing pCGN10822, which provides of the expression of the ATPT2 sequence,  
10 in the sense orientation, from the napin promoter. Provided are graphs for alpha, gamma, and delta tocopherols, as well as total tocopherol for 22 transformed lines, as well as a nontransformed (wildtype) control.

Figure 25 provides a bar graph of HPLC analysis of seed extracts from *Arabidopsis* plants transformed with pCGN10803 (35S-ATPT2, in the antisense orientation), pCGN10822 (line  
15 1625, napin ATPT2 in the sense orientation), pCGN10809 (line 1627, 35S-ATPT3 in the sense orientation), a nontransformed (wt) control, and an empty vector transformed control.

Figure 26 shows total tocopherol levels measured in T# *Arabidopsis* seed of line.

Figure 27 shows total tocopherol levels measured in T# *Arabidopsis* seed of line.

Figure 28 shows total tocopherol levels measured in developing canola seed of line  
20 10822-1.

Figure 29: shows results of phytyl prenyltransferase activity assay using *Synechocystis* wild type and slr1737 knockout mutant membrane preparations.

Figure 30 is the chromatograph from an HPLC analysis of *Synechocystis* extracts.

Figure 31 is a sequence alignment of the *Arabidopsis* homologue with the sequence of the  
25 public database.

Figure 32 shows the results of hydropathic analysis of slr1737

Figure 33 shows the results of hydropathic analysis of the *Arabidopsis* homologue of slr1737.

Figure 34 shows the catalytic mechanism of various cyclase enzymes

Figure 35 is a sequence alignment of slr1737, slr1737 *Arabidopsis* homologue and the *Arabidopsis* chalcone isomerase.

## DETAILED DESCRIPTION OF THE INVENTION

5

The present invention provides, *inter alia*, compositions and methods for altering (for example, increasing and decreasing) the tocopherol levels and/or modulating their ratios in host cells. In particular, the present invention provides polynucleotides, polypeptides, and methods of use thereof for the modulation of tocopherol content in host plant cells.

10

The biosynthesis of  $\alpha$ -tocopherol in higher plants involves condensation of homogentisic acid and phytylpyrophosphate to form 2-methyl-6 phytylbenzoquinol that can, by cyclization and subsequent methylations (Fiedler et al., 1982, *Planta*, 155: 511-515, Soll et al., 1980, *Arch. Biochem. Biophys.* 204: 544-550, Marshall et al., 1985 *Phytochem.*, 24: 1705-1711, all of which are herein incorporated by reference in their entirety), form various tocopherols.

15

The *Arabidopsis pds2* mutant identified and characterized by Norris *et al.* (1995), is deficient in tocopherol and plastiquinone-9 accumulation. Further genetic and biochemical analysis suggested that the protein encoded by *PDS2* may be responsible for the prenylation of homogentisic acid. The *PDS2* locus identified by Norris *et al.* (1995) has been hypothesized to possibly encode the tocopherol phytyl-prenyltransferase, as the *pds2* mutant fails to accumulate

20

tocopherols.

25

Norris *et al.* (1995) determined that in *Arabidopsis pds2* lies at the top of chromosome 3, approximately 7 centimorgans above long hypocotyl2, based on the genetic map. ATPT2 is located on chromosome 2 between 36 and 41 centimorgans, lying on BAC F19F24, indicating that ATPT2 does not correspond to *PDS2*. Thus, it is an aspect of the present invention to provide novel polynucleotides and polypeptides involved in the prenylation of homogentisic acid. This reaction may be a rate limiting step in tocopherol biosynthesis, and this gene has yet to be isolated.

U.S. Patent No. 5,432,069 describes the partial purification and characterization of tocopherol cyclase from *Chlorella protothecoides*, *Dunaliella salina* and wheat. The cyclase

described as being glycine rich, water soluble and with a predicted MW of 48-50kDa. However, only limited peptide fragment sequences were available.

In one aspect, the present invention provides polynucleotide and polypeptide sequences involved in the prenylation of straight chain and aromatic compounds. Straight chain prenyltransferases as used herein comprises sequences which encode proteins involved in the prenylation of straight chain compounds, including, but not limited to, geranyl geranyl pyrophosphate and farnesyl pyrophosphate. Aromatic prenyltransferases, as used herein, comprises sequences which encode proteins involved in the prenylation of aromatic compounds, including, but not limited to, menaquinone, ubiquinone, chlorophyll, and homogentisic acid. The prenyltransferase of the present invention preferably prenylates homogentisic acid.

In another aspect, the invention provides polynucleotide and polypeptide sequences to tocopherol cyclization enzymes. The 2,3-dimethyl-5-phytylplastoquinol cyclase (tocopherol cyclase) is responsible for the cyclization of 2,3-dimethyl-5-phytylplastoquinol to tocopherol.

#### 15 Isolated Polynucleotides, Proteins, and Polypeptides

A first aspect of the present invention relates to isolated prenyltransferase polynucleotides. Another aspect of the present invention relates to isolated tocopherol cyclase polynucleotides. The polynucleotide sequences of the present invention include isolated polynucleotides that encode the polypeptides of the invention having a deduced amino acid sequence selected from the group of sequences set forth in the Sequence Listing and to other polynucleotide sequences closely related to such sequences and variants thereof.

The invention provides a polynucleotide sequence identical over its entire length to each coding sequence as set forth in the Sequence Listing. The invention also provides the coding sequence for the mature polypeptide or a fragment thereof, as well as the coding sequence for the mature polypeptide or a fragment thereof in a reading frame with other coding sequences, such as those encoding a leader or secretory sequence, a pre-, pro-, or prepro- protein sequence. The polynucleotide can also include non-coding sequences, including for example, but not limited to, non-coding 5' and 3' sequences, such as the transcribed, untranslated sequences, termination signals, ribosome binding sites, sequences that stabilize mRNA, introns, polyadenylation signals,

and additional coding sequence that encodes additional amino acids. For example, a marker sequence can be included to facilitate the purification of the fused polypeptide. Polynucleotides of the present invention also include polynucleotides comprising a structural gene and the naturally associated sequences that control gene expression.

5 The invention also includes polynucleotides of the formula:



wherein, at the 5' end, X is hydrogen, and at the 3' end, Y is hydrogen or a metal,  $R_1$  and  $R_3$  are any nucleic acid residue,  $n$  is an integer between 1 and 3000, preferably between 1 and 1000 and  $R_2$  is a nucleic acid sequence of the invention, particularly a nucleic acid sequence selected from  
10 the group set forth in the Sequence Listing and preferably those of SEQ ID NOs: 1, 3, 5, 7, 8, 10, 11, 13-16, 18, 23, 29, 36, and 38. In the formula,  $R_2$  is oriented so that its 5' end residue is at the left, bound to  $R_1$ , and its 3' end residue is at the right, bound to  $R_3$ . Any stretch of nucleic acid residues denoted by either R group, where R is greater than 1, may be either a heteropolymer or a homopolymer, preferably a heteropolymer.

15 The invention also relates to variants of the polynucleotides described herein that encode for variants of the polypeptides of the invention. Variants that are fragments of the polynucleotides of the invention can be used to synthesize full-length polynucleotides of the invention. Preferred embodiments are polynucleotides encoding polypeptide variants wherein 5 to 10, 1 to 5, 1 to 3, 2, 1 or no amino acid residues of a polypeptide sequence of the invention are  
20 substituted, added or deleted, in any combination. Particularly preferred are substitutions, additions, and deletions that are silent such that they do not alter the properties or activities of the polynucleotide or polypeptide.

Further preferred embodiments of the invention that are at least 50%, 60%, or 70% identical over their entire length to a polynucleotide encoding a polypeptide of the invention, and  
25 polynucleotides that are complementary to such polynucleotides. More preferable are polynucleotides that comprise a region that is at least 80% identical over its entire length to a polynucleotide encoding a polypeptide of the invention and polynucleotides that are complementary thereto. In this regard, polynucleotides at least 90% identical over their entire length are particularly preferred, those at least 95% identical are especially preferred. Further,



those with at least 97% identity are highly preferred and those with at least 98% and 99% identity are particularly highly preferred, with those at least 99% being the most highly preferred.

Preferred embodiments are polynucleotides that encode polypeptides that retain substantially the same biological function or activity as the mature polypeptides encoded by the polynucleotides set forth in the Sequence Listing.

The invention further relates to polynucleotides that hybridize to the above-described sequences. In particular, the invention relates to polynucleotides that hybridize under stringent conditions to the above-described polynucleotides. As used herein, the terms "stringent conditions" and "stringent hybridization conditions" mean that hybridization will generally occur if there is at least 95% and preferably at least 97% identity between the sequences. An example of stringent hybridization conditions is overnight incubation at 42°C in a solution comprising 50% formamide, 5x SSC (150 mM NaCl, 15 mM trisodium citrate), 50 mM sodium phosphate (pH 7.6), 5x Denhardt's solution, 10% dextran sulfate, and 20 micrograms/milliliter denatured, sheared salmon sperm DNA, followed by washing the hybridization support in 0.1x SSC at approximately 65°C. Other hybridization and wash conditions are well known and are exemplified in Sambrook, *et al.*, Molecular Cloning: A Laboratory Manual, Second Edition, Cold Spring Harbor, NY (1989), particularly Chapter 11.

The invention also provides a polynucleotide consisting essentially of a polynucleotide sequence obtainable by screening an appropriate library containing the complete gene for a polynucleotide sequence set forth in the Sequence Listing under stringent hybridization conditions with a probe having the sequence of said polynucleotide sequence or a fragment thereof; and isolating said polynucleotide sequence. Fragments useful for obtaining such a polynucleotide include, for example, probes and primers as described herein.

As discussed herein regarding polynucleotide assays of the invention, for example, polynucleotides of the invention can be used as a hybridization probe for RNA, cDNA, or genomic DNA to isolate full length cDNAs or genomic clones encoding a polypeptide and to isolate cDNA or genomic clones of other genes that have a high sequence similarity to a polynucleotide set forth in the Sequence Listing. Such probes will generally comprise at least 15 bases. Preferably such probes will have at least 30 bases and can have at least 50 bases. Particularly preferred probes will have between 30 bases and 50 bases, inclusive.

The coding region of each gene that comprises or is comprised by a polynucleotide sequence set forth in the Sequence Listing may be isolated by screening using a DNA sequence provided in the Sequence Listing to synthesize an oligonucleotide probe. A labeled oligonucleotide having a sequence complementary to that of a gene of the invention is then used to screen a library of cDNA, genomic DNA or mRNA to identify members of the library which hybridize to the probe. For example, synthetic oligonucleotides are prepared which correspond to the prenyltransferase or tocopherol cyclase EST sequences. The oligonucleotides are used as primers in polymerase chain reaction (PCR) techniques to obtain 5' and 3' terminal sequence of prenyltransferase or tocopherol cyclase genes. Alternatively, where oligonucleotides of low degeneracy can be prepared from particular prenyltransferase or tocopherol cyclase peptides, such probes may be used directly to screen gene libraries for prenyltransferase or tocopherol cyclase gene sequences. In particular, screening of cDNA libraries in phage vectors is useful in such methods due to lower levels of background hybridization.

Typically, a prenyltransferase or tocopherol cyclase sequence obtainable from the use of nucleic acid probes will show 60-70% sequence identity between the target prenyltransferase or tocopherol cyclase sequence and the encoding sequence used as a probe. However, lengthy sequences with as little as 50-60% sequence identity may also be obtained. The nucleic acid probes may be a lengthy fragment of the nucleic acid sequence, or may also be a shorter, oligonucleotide probe. When longer nucleic acid fragments are employed as probes (greater than about 100 bp), one may screen at lower stringencies in order to obtain sequences from the target sample which have 20-50% deviation (i.e., 50-80% sequence homology) from the sequences used as probe. Oligonucleotide probes can be considerably shorter than the entire nucleic acid sequence encoding an prenyltransferase or tocopherol cyclase enzyme, but should be at least about 10, preferably at least about 15, and more preferably at least about 20 nucleotides. A higher degree of sequence identity is desired when shorter regions are used as opposed to longer regions. It may thus be desirable to identify regions of highly conserved amino acid sequence to design oligonucleotide probes for detecting and recovering other related prenyltransferase or tocopherol cyclase genes. Shorter probes are often particularly useful for polymerase chain reactions (PCR), especially when highly conserved sequences can be identified. (See, Gould, *et al.*, *PNAS USA* (1989) 86:1934-1938.).

Another aspect of the present invention relates to prenyltransferase or tocopherol cyclase polypeptides. Such polypeptides include isolated polypeptides set forth in the Sequence Listing, as well as polypeptides and fragments thereof, particularly those polypeptides which exhibit prenyltransferase or tocopherol cyclase activity and also those polypeptides which have at least 50%, 60% or 70% identity, preferably at least 80% identity, more preferably at least 90% identity, and most preferably at least 95% identity to a polypeptide sequence selected from the group of sequences set forth in the Sequence Listing, and also include portions of such polypeptides, wherein such portion of the polypeptide preferably includes at least 30 amino acids and more preferably includes at least 50 amino acids.

“Identity”, as is well understood in the art, is a relationship between two or more polypeptide sequences or two or more polynucleotide sequences, as determined by comparing the sequences. In the art, “identity” also means the degree of sequence relatedness between polypeptide or polynucleotide sequences, as determined by the match between strings of such sequences. “Identity” can be readily calculated by known methods including, but not limited to, those described in *Computational Molecular Biology*, Lesk, A.M., ed., Oxford University Press, New York (1988); *Biocomputing: Informatics and Genome Projects*, Smith, D.W., ed., Academic Press, New York, 1993; *Computer Analysis of Sequence Data, Part I*, Griffin, A.M. and Griffin, H.G., eds., Humana Press, New Jersey (1994); *Sequence Analysis in Molecular Biology*, von Heinje, G., Academic Press (1987); *Sequence Analysis Primer*, Gribskov, M. and Devereux, J., eds., Stockton Press, New York (1991); and Carillo, H., and Lipman, D., *SIAM J Applied Math*, 48:1073 (1988). Methods to determine identity are designed to give the largest match between the sequences tested. Moreover, methods to determine identity are codified in publicly available programs. Computer programs which can be used to determine identity between two sequences include, but are not limited to, GCG (Devereux, J., et al., *Nucleic Acids Research* 12(1):387 (1984); suite of five BLAST programs, three designed for nucleotide sequences queries (BLASTN, BLASTX, and TBLASTX) and two designed for protein sequence queries (BLASTP and TBLASTN) (Coulson, *Trends in Biotechnology*, 12: 76-80 (1994); Birren, et al., *Genome Analysis*, 1: 543-559 (1997)). The BLAST X program is publicly available from NCBI and other sources (*BLAST Manual*, Altschul, S., et al., NCBI NLM NIH, Bethesda, MD

20894; Altschul, S., *et al.*, *J. Mol. Biol.*, 215:403-410 (1990)). The well known Smith Waterman algorithm can also be used to determine identity.

Parameters for polypeptide sequence comparison typically include the following:

Algorithm: Needleman and Wunsch, *J. Mol. Biol.* 48:443-453 (1970)

5 Comparison matrix: BLOSSUM62 from Hentikoff and Hentikoff, *Proc. Natl. Acad. Sci USA* 89:10915-10919 (1992)

Gap Penalty: 12

Gap Length Penalty: 4

10 A program which can be used with these parameters is publicly available as the "gap" program from Genetics Computer Group, Madison Wisconsin. The above parameters along with no penalty for end gap are the default parameters for peptide comparisons.

Parameters for polynucleotide sequence comparison include the following:

Algorithm: Needleman and Wunsch, *J. Mol. Biol.* 48:443-453 (1970)

Comparison matrix: matches = +10; mismatches = 0

15 Gap Penalty: 50

Gap Length Penalty: 3

A program which can be used with these parameters is publicly available as the "gap" program from Genetics Computer Group, Madison Wisconsin. The above parameters are the default parameters for nucleic acid comparisons.

20 The invention also includes polypeptides of the formula:



wherein, at the amino terminus, X is hydrogen, and at the carboxyl terminus, Y is hydrogen or a metal,  $R_1$  and  $R_3$  are any amino acid residue,  $n$  is an integer between 1 and 1000, and  $R_2$  is an amino acid sequence of the invention, particularly an amino acid sequence selected from the group set forth in the Sequence Listing and preferably those encoded by the sequences provided in SEQ ID NOs: 2, 4, 6, 9, 12, 17, 19-22, 24-28, 30, 32-35, 37, and 39. In the formula,  $R_2$  is oriented so that its amino terminal residue is at the left, bound to  $R_1$ , and its carboxy terminal residue is at the right, bound to  $R_3$ . Any stretch of amino acid residues denoted by either R group, where R is greater than 1, may be either a heteropolymer or a homopolymer, preferably a heteropolymer.

25

30

Polypeptides of the present invention include isolated polypeptides encoded by a polynucleotide comprising a sequence selected from the group of a sequence contained in the Sequence Listing set forth herein.

5 The polypeptides of the present invention can be mature protein or can be part of a fusion protein.

Fragments and variants of the polypeptides are also considered to be a part of the invention. A fragment is a variant polypeptide which has an amino acid sequence that is entirely the same as part but not all of the amino acid sequence of the previously described polypeptides. The fragments can be "free-standing" or comprised within a larger polypeptide of which the  
10 fragment forms a part or a region, most preferably as a single continuous region. Preferred fragments are biologically active fragments which are those fragments that mediate activities of the polypeptides of the invention, including those with similar activity or improved activity or with a decreased activity. Also included are those fragments that antigenic or immunogenic in an animal, particularly a human.

15 Variants of the polypeptide also include polypeptides that vary from the sequences set forth in the Sequence Listing by conservative amino acid substitutions, substitution of a residue by another with like characteristics. In general, such substitutions are among Ala, Val, Leu and Ile; between Ser and Thr; between Asp and Glu; between Asn and Gln; between Lys and Arg; or between Phe and Tyr. Particularly preferred are variants in which 5 to 10; 1 to 5; 1 to 3 or one  
20 amino acid(s) are substituted, deleted, or added, in any combination.

Variants that are fragments of the polypeptides of the invention can be used to produce the corresponding full length polypeptide by peptide synthesis. Therefore, these variants can be used as intermediates for producing the full-length polypeptides of the invention.

25 The polynucleotides and polypeptides of the invention can be used, for example, in the transformation of host cells, such as plant host cells, as further discussed herein.

The invention also provides polynucleotides that encode a polypeptide that is a mature protein plus additional amino or carboxyl-terminal amino acids, or amino acids within the mature polypeptide (for example, when the mature form of the protein has more than one polypeptide chain). Such sequences can, for example, play a role in the processing of a protein from a  
30 precursor to a mature form, allow protein transport, shorten or lengthen protein half-life, or



facilitate manipulation of the protein in assays or production. It is contemplated that cellular enzymes can be used to remove any additional amino acids from the mature protein.

A precursor protein, having the mature form of the polypeptide fused to one or more prosequences may be an inactive form of the polypeptide. The inactive precursors generally are activated when the prosequences are removed. Some or all of the prosequences may be removed prior to activation. Such precursor protein are generally called proproteins.

### Plant Constructs and Methods of Use

Of particular interest is the use of the nucleotide sequences in recombinant DNA constructs to direct the transcription or transcription and translation (expression) of the prenyltransferase or tocopherol cyclase sequences of the present invention in a host plant cell. The expression constructs generally comprise a promoter functional in a host plant cell operably linked to a nucleic acid sequence encoding a prenyltransferase or tocopherol cyclase of the present invention and a transcriptional termination region functional in a host plant cell.

A first nucleic acid sequence is "operably linked" or "operably associated" with a second nucleic acid sequence when the sequences are so arranged that the first nucleic acid sequence affects the function of the second nucleic-acid sequence. Preferably, the two sequences are part of a single contiguous nucleic acid molecule and more preferably are adjacent. For example, a promoter is operably linked to a gene if the promoter regulates or mediates transcription of the gene in a cell.

Those skilled in the art will recognize that there are a number of promoters which are functional in plant cells, and have been described in the literature. Chloroplast and plastid specific promoters, chloroplast or plastid functional promoters, and chloroplast or plastid operable promoters are also envisioned.

One set of plant functional promoters are constitutive promoters such as the CaMV35S or FMV35S promoters that yield high levels of expression in most plant organs. Enhanced or duplicated versions of the CaMV35S and FMV35S promoters are useful in the practice of this invention (Odell, *et al.* (1985) *Nature* 313:810-812; Rogers, U.S. Patent Number 5,378, 619). In addition, it may also be preferred to bring about expression of the prenyltransferase or tocopherol

cyclase gene in specific tissues of the plant, such as leaf, stem, root, tuber, seed, fruit, etc., and the promoter chosen should have the desired tissue and developmental specificity.

Of particular interest is the expression of the nucleic acid sequences of the present invention from transcription initiation regions which are preferentially expressed in a plant seed  
5 tissue. Examples of such seed preferential transcription initiation sequences include those sequences derived from sequences encoding plant storage protein genes or from genes involved in fatty acid biosynthesis in oilseeds. Examples of such promoters include the 5' regulatory regions from such genes as napin (Kridl *et al.*, *Seed Sci. Res.* 1:209:219 (1991)), phaseolin, zein, soybean trypsin inhibitor, ACP, stearyl-ACP desaturase, soybean  $\alpha'$  subunit of  $\beta$ -conglycinin  
10 (soy 7s, (Chen *et al.*, *Proc. Natl. Acad. Sci.*, 83:8560-8564 (1986))) and oleosin.

It may be advantageous to direct the localization of proteins conferring prenyltransferase or tocopherol cyclase to a particular subcellular compartment, for example, to the mitochondrion, endoplasmic reticulum, vacuoles, chloroplast or other plastidic compartment. For example, where the genes of interest of the present invention will be targeted to plastids, such as  
15 chloroplasts, for expression, the constructs will also employ the use of sequences to direct the gene to the plastid. Such sequences are referred to herein as chloroplast transit peptides (CTP) or plastid transit peptides (PTP). In this manner, where the gene of interest is not directly inserted into the plastid, the expression construct will additionally contain a gene encoding a transit peptide to direct the gene of interest to the plastid. The chloroplast transit peptides may be  
20 derived from the gene of interest, or may be derived from a heterologous sequence having a CTP. Such transit peptides are known in the art. See, for example, Von Heijne *et al.* (1991) *Plant Mol. Biol. Rep.* 9:104-126; Clark *et al.* (1989) *J. Biol. Chem.* 264:17544-17550; della-Cioppa *et al.* (1987) *Plant Physiol.* 84:965-968; Romer *et al.* (1993) *Biochem. Biophys. Res Commun.* 196:1414-1421; and, Shah *et al.* (1986) *Science* 233:478-481.

25 Depending upon the intended use, the constructs may contain the nucleic acid sequence which encodes the entire prenyltransferase or tocopherol cyclase protein, or a portion thereof. For example, where antisense inhibition of a given prenyltransferase or tocopherol cyclase protein is desired, the entire prenyltransferase or tocopherol cyclase sequence is not required. Furthermore, where prenyltransferase or tocopherol cyclase sequences used in constructs are  
30 intended for use as probes, it may be advantageous to prepare constructs containing only a

particular portion of a prenyltransferase or tocopherol cyclase encoding sequence, for example a sequence which is discovered to encode a highly conserved prenyltransferase or tocopherol cyclase region.

The skilled artisan will recognize that there are various methods for the inhibition of expression of endogenous sequences in a host cell. Such methods include, but are not limited to, antisense suppression (Smith, *et al.* (1988) *Nature* 334:724-726), co-suppression (Napoli, *et al.* (1989) *Plant Cell* 2:279-289), ribozymes (PCT Publication WO 97/10328), and combinations of sense and antisense Waterhouse, *et al.* (1998) *Proc. Natl. Acad. Sci. USA* 95:13959-13964. Methods for the suppression of endogenous sequences in a host cell typically employ the transcription or transcription and translation of at least a portion of the sequence to be suppressed. Such sequences may be homologous to coding as well as non-coding regions of the endogenous sequence.

Regulatory transcript termination regions may be provided in plant expression constructs of this invention as well. Transcript termination regions may be provided by the DNA sequence encoding the prenyltransferase or tocopherol cyclase or a convenient transcription termination region derived from a different gene source, for example, the transcript termination region which is naturally associated with the transcript initiation region. The skilled artisan will recognize that any convenient transcript termination region which is capable of terminating transcription in a plant cell may be employed in the constructs of the present invention.

Alternatively, constructs may be prepared to direct the expression of the prenyltransferase or tocopherol cyclase sequences directly from the host plant cell plastid. Such constructs and methods are known in the art and are generally described, for example, in Svab, *et al.* (1990) *Proc. Natl. Acad. Sci. USA* 87:8526-8530 and Svab and Maliga (1993) *Proc. Natl. Acad. Sci. USA* 90:913-917 and in U.S. Patent Number 5,693,507.

The prenyltransferase or tocopherol cyclase constructs of the present invention can be used in transformation methods with additional constructs providing for the expression of other nucleic acid sequences encoding proteins involved in the production of tocopherols, or tocopherol precursors such as homogentisic acid and/or phytylpyrophosphate. Nucleic acid sequences encoding proteins involved in the production of homogentisic acid are known in the art, and include but are not limited to, 4-hydroxyphenylpyruvate dioxygenase (HPPD, EC

1.13.11.27) described for example, by Garcia, *et al.* ((1999) *Plant Physiol.* 119(4):1507-1516), mono or bifunctional tyrA (described for example by Xia, *et al.* (1992) *J. Gen Microbiol.* 138:1309-1316, and Hudson, *et al.* (1984) *J. Mol. Biol.* 180:1023-1051), Oxygenase, 4-hydroxyphenylpyruvate di- (9Cl), 4-Hydroxyphenylpyruvate dioxygenase; p-Hydroxyphenylpyruvate dioxygenase; p-Hydroxyphenylpyruvate hydroxylase; p-Hydroxyphenylpyruvate oxidase; p-Hydroxyphenylpyruvic acid hydroxylase; p-Hydroxyphenylpyruvic hydroxylase; p-Hydroxyphenylpyruvic oxidase), 4-hydroxyphenylacetate, NAD(P)H:oxygen oxidoreductase (1-hydroxylating); 4-hydroxyphenylacetate 1-monooxygenase, and the like. In addition, constructs for the expression of nucleic acid sequences encoding proteins involved in the production of phytylpyrophosphate can also be employed with the prenyltransferase or tocopherol cyclase constructs of the present invention. Nucleic acid sequences encoding proteins involved in the production of phytylpyrophosphate are known in the art, and include, but are not limited to geranylgeranylpyrophosphate synthase (GGPPS), geranylgeranylpyrophosphate reductase (GGH), 1-deoxyxylulose-5-phosphate synthase, 1-deoxy-D-xylulose-5-phosphate reductoisomerase, 4-diphosphocytidyl-2-C-methylerythritol synthase, isopentyl pyrophosphate isomerase.

The prenyltransferase or tocopherol cyclase sequences of the present invention find use in the preparation of transformation constructs having a second expression cassette for the expression of additional sequences involved in tocopherol biosynthesis. Additional tocopherol biosynthesis sequences of interest in the present invention include, but are not limited to gamma-tocopherol methyltransferase (Shintani, *et al.* (1998) *Science* 282(5396):2098-2100), tocopherol cyclase, and tocopherol methyltransferase.

A plant cell, tissue, organ, or plant into which the recombinant DNA constructs containing the expression constructs have been introduced is considered transformed, transfected, or transgenic. A transgenic or transformed cell or plant also includes progeny of the cell or plant and progeny produced from a breeding program employing such a transgenic plant as a parent in a cross and exhibiting an altered phenotype resulting from the presence of a prenyltransferase or tocopherol cyclase nucleic acid sequence.

Plant expression or transcription constructs having a prenyltransferase or tocopherol cyclase as the DNA sequence of interest for increased or decreased expression thereof may be

employed with a wide variety of plant life. particularly. plant life involved in the production of vegetable oils for edible and industrial uses. Particularly preferred plants for use in the methods of the present invention include, but are not limited to: *Acacia*, alfalfa, aneth, apple, apricot, artichoke, arugula, asparagus, avocado, banana, barley, beans, beet, blackberry, blueberry, broccoli, brussels sprouts, cabbage, canola, cantaloupe, carrot, cassava, cauliflower, celery, 5 cherry, chicory, cilantro, citrus, clementines, coffee, corn, cotton, cucumber, Douglas fir, eggplant, endive, escarole, eucalyptus, fennel, figs, garlic, gourd, grape, grapefruit, honey dew, jicama, kiwifruit, lettuce, leeks, lemon, lime, Loblolly pine, mango, melon, mushroom, nectarine, nut, oat, oil palm, oil seed rape, okra, onion, orange, an ornamental plant, papaya, parsley, pea, 10 peach, peanut, pear, pepper, persimmon, pine, pineapple, plantain, plum, pomegranate, poplar, potato, pumpkin, quince, radiata pine, radicchio, radish, raspberry, rice, rye, sorghum, Southern pine, soybean, spinach, squash, strawberry, sugarbeet, sugarcane, sunflower, sweet potato, sweetgum, tangerine, tea, tobacco, tomato, triticale, turf, turnip, a vine, watermelon, wheat, yams, and zucchini.

15 . . . . . Most especially preferred are temperate oilseed crops. Temperate oilseed crops of interest include, but are not limited to, rapeseed (Canola and High Erucic Acid varieties), sunflower, safflower, cotton, soybean, peanut, coconut and oil palms, and corn. Depending on the method for introducing the recombinant constructs into the host cell, other DNA sequences may be required. Importantly, this invention is applicable to dicotyledons and monocotyledons 20 species alike and will be readily applicable to new and/or improved transformation and regulation techniques.

Of particular interest, is the use of prenyltransferase or tocopherol cyclase constructs in plants to produce plants or plant parts, including, but not limited to leaves, stems, roots, reproductive, and seed, with a modified content of tocopherols in plant parts having transformed 25 plant cells.

For immunological screening, antibodies to the protein can be prepared by injecting rabbits or mice with the purified protein or portion thereof, such methods of preparing antibodies being well known to those in the art. Either monoclonal or polyclonal antibodies can be produced, although typically polyclonal antibodies are more useful for gene isolation. Western 30 analysis may be conducted to determine that a related protein is present in a crude extract of the



desired plant species, as determined by cross-reaction with the antibodies to the encoded proteins. When cross-reactivity is observed, genes encoding the related proteins are isolated by screening expression libraries representing the desired plant species. Expression libraries can be constructed in a variety of commercially available vectors, including lambda gt11, as described in  
5 Sambrook, *et al.* (*Molecular Cloning: A Laboratory Manual*, Second Edition (1989) Cold Spring Harbor Laboratory, Cold Spring Harbor, New York).

To confirm the activity and specificity of the proteins encoded by the identified nucleic acid sequences as prenyltransferase or tocopherol cyclase enzymes, *in vitro* assays are performed in insect cell cultures using baculovirus expression systems. Such baculovirus expression  
10 systems are known in the art and are described by Lee, *et al.* U.S. Patent Number 5,348,886, the entirety of which is herein incorporated by reference.

In addition, other expression constructs may be prepared to assay for protein activity utilizing different expression systems. Such expression constructs are transformed into yeast or prokaryotic host and assayed for prenyltransferase or tocopherol cyclase activity. Such  
15 expression systems are known in the art and are readily available through commercial sources.

In addition to the sequences described in the present invention, DNA coding sequences useful in the present invention can be derived from algae, fungi, bacteria, mammalian sources, plants, etc. Homology searches in existing databases using signature sequences corresponding to conserved nucleotide and amino acid sequences of prenyltransferase or tocopherol cyclase can  
20 be employed to isolate equivalent, related genes from other sources such as plants and microorganisms. Searches in EST databases can also be employed. Furthermore, the use of DNA sequences encoding enzymes functionally enzymatically equivalent to those disclosed herein, wherein such DNA sequences are degenerate equivalents of the nucleic acid sequences disclosed herein in accordance with the degeneracy of the genetic code, is also encompassed by  
25 the present invention. Demonstration of the functionality of coding sequences identified by any of these methods can be carried out by complementation of mutants of appropriate organisms, such as *Synechocystis*, *Shewanella*, yeast, *Pseudomonas*, *Rhodobacteria*, etc., that lack specific biochemical reactions, or that have been mutated. The sequences of the DNA coding regions can be optimized by gene resynthesis, based on codon usage, for maximum expression in  
30 particular hosts.

For the alteration of tocopherol production in a host cell, a second expression construct can be used in accordance with the present invention. For example, the prenyltransferase or tocopherol cyclase expression construct can be introduced into a host cell in conjunction with a second expression construct having a nucleotide sequence for a protein involved in tocopherol biosynthesis.

The method of transformation in obtaining such transgenic plants is not critical to the instant invention, and various methods of plant transformation are currently available. Furthermore, as newer methods become available to transform crops, they may also be directly applied hereunder. For example, many plant species naturally susceptible to *Agrobacterium* infection may be successfully transformed via tripartite or binary vector methods of *Agrobacterium* mediated transformation. In many instances, it will be desirable to have the construct bordered on one or both sides by T-DNA, particularly having the left and right borders, more particularly the right border. This is particularly useful when the construct uses *A. tumefaciens* or *A. rhizogenes* as a mode for transformation, although the T-DNA borders may find use with other modes of transformation. In addition, techniques of microinjection, DNA particle bombardment, and electroporation have been developed which allow for the transformation of various monocot and dicot plant species.

Normally, included with the DNA construct will be a structural gene having the necessary regulatory regions for expression in a host and providing for selection of transformant cells. The gene may provide for resistance to a cytotoxic agent, e.g. antibiotic, heavy metal, toxin, etc., complementation providing prototrophy to an auxotrophic host, viral immunity or the like. Depending upon the number of different host species the expression construct or components thereof are introduced, one or more markers may be employed, where different conditions for selection are used for the different hosts.

Where *Agrobacterium* is used for plant cell transformation, a vector may be used which may be introduced into the *Agrobacterium* host for homologous recombination with T-DNA or the Ti- or Ri-plasmid present in the *Agrobacterium* host. The Ti- or Ri-plasmid containing the T-DNA for recombination may be armed (capable of causing gall formation) or disarmed (incapable of causing gall formation), the latter being permissible, so long as the *vir* genes are

present in the transformed *Agrobacterium* host. The armed plasmid can give a mixture of normal plant cells and gall.

In some instances where *Agrobacterium* is used as the vehicle for transforming host plant cells, the expression or transcription construct bordered by the T-DNA border region(s) will be inserted into a broad host range vector capable of replication in *E. coli* and *Agrobacterium*, there being broad host range vectors described in the literature. Commonly used is pRK2 or derivatives thereof. See, for example, Ditta, *et al.*, (*Proc. Nat. Acad. Sci., U.S.A.* (1980) 77:7347-7351) and EPA 0 120 515, which are incorporated herein by reference. Alternatively, one may insert the sequences to be expressed in plant cells into a vector containing separate replication sequences, one of which stabilizes the vector in *E. coli*, and the other in *Agrobacterium*. See, for example, McBride, *et al.* (*Plant Mol. Biol.* (1990) 14:269-276), wherein the pRiHRI (Jouanin, *et al.*, *Mol. Gen. Genet.* (1985) 201:370-374) origin of replication is utilized and provides for added stability of the plant expression vectors in host *Agrobacterium* cells.

Included with the expression construct and the T-DNA will be one or more markers, which allow for selection of transformed *Agrobacterium* and transformed plant cells. A number of markers have been developed for use with plant cells, such as resistance to chloramphenicol, kanamycin, the aminoglycoside G418, hygromycin, or the like. The particular marker employed is not essential to this invention, one or another marker being preferred depending on the particular host and the manner of construction.

For transformation of plant cells using *Agrobacterium*, explants may be combined and incubated with the transformed *Agrobacterium* for sufficient time for transformation, the bacteria killed, and the plant cells cultured in an appropriate selective medium. Once callus forms, shoot formation can be encouraged by employing the appropriate plant hormones in accordance with known methods and the shoots transferred to rooting medium for regeneration of plants. The plants may then be grown to seed and the seed used to establish repetitive generations and for isolation of vegetable oils.

There are several possible ways to obtain the plant cells of this invention which contain multiple expression constructs. Any means for producing a plant comprising a construct having a DNA sequence encoding the expression construct of the present invention, and at least one

other construct having another DNA sequence encoding an enzyme are encompassed by the present invention. For example, the expression construct can be used to transform a plant at the same time as the second construct either by inclusion of both expression constructs in a single transformation vector or by using separate vectors, each of which express desired genes. The  
5 second construct can be introduced into a plant which has already been transformed with the prenyltransferase or tocopherol cyclase expression construct, or alternatively, transformed plants, one expressing the prenyltransferase or tocopherol cyclase construct and one expressing the second construct, can be crossed to bring the constructs together in the same plant.

Transgenic plants of the present invention may be produced from tissue culture, and  
10 subsequent generations grown from seed. Alternatively, transgenic plants may be grown using apomixis. Apomixis is a genetically controlled method of reproduction in plants where the embryo is formed without union of an egg and a sperm. There are three basic types of apomictic reproduction: 1) apospory where the embryo develops from a chromosomally unreduced egg in an embryo sac derived from the nucleus, 2) diplospory where the embryo develops from an  
15 unreduced egg in an embryo sac derived from the megaspore mother cell, and 3) adventitious embryony where the embryo develops directly from a somatic cell. In most forms of apomixis, pseudogamy or fertilization of the polar nuclei to produce endosperm is necessary for seed viability. In apospory, a nurse cultivar can be used as a pollen source for endosperm formation in seeds. The nurse cultivar does not affect the genetics of the aposporous apomictic cultivar since  
20 the unreduced egg of the cultivar develops parthenogenetically, but makes possible endosperm production. Apomixis is economically important, especially in transgenic plants, because it causes any genotype, no matter how heterozygous, to breed true. Thus, with apomictic reproduction, heterozygous transgenic plants can maintain their genetic fidelity throughout repeated life cycles. Methods for the production of apomictic plants are known in the art. See,  
25 U.S. Patent No. 5,811,636, which is herein incorporated by reference in its entirety.

The nucleic acid sequences of the present invention can be used in constructs to provide for the expression of the sequence in a variety of host cells, both prokaryotic eukaryotic. Host cells of the present invention preferably include monocotyledenous and dicotyledenous plant cells.

In general, the skilled artisan is familiar with the standard resource materials which describe specific conditions and procedures for the construction, manipulation and isolation of macromolecules (e.g., DNA molecules, plasmids, etc.), generation of recombinant organisms and the screening and isolating of clones, (see for example, Sambrook *et al.*, *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Press (1989); Maliga *et al.*, *Methods in Plant Molecular Biology*, Cold Spring Harbor Press (1995), the entirety of which is herein incorporated by reference; Birren *et al.*, *Genome Analysis: Analyzing DNA*, 1, Cold Spring Harbor, New York, the entirety of which is herein incorporated by reference).

Methods for the expression of sequences in insect host cells are known in the art.

10 Baculovirus expression vectors are recombinant insect viruses in which the coding sequence for a chosen foreign gene has been inserted behind a baculovirus promoter in place of the viral gene, e.g., polyhedrin (Smith and Summers, U.S. Pat. No., 4,745,051, the entirety of which is incorporated herein by reference). Baculovirus expression vectors are known in the art, and are described for example in Doerfler, *Curr. Top. Microbiol. Immunol.* 131:51-68 (1968); Luckow  
15 and Summers, *Bio/Technology* 6:47-55 (1988a); Miller, *Annual Review of Microbiol.* 42:177-199 (1988); Summers, *Curr. Comm. Molecular Biology*, Cold Spring Harbor Press, Cold Spring Harbor, N.Y. (1988); Summers and Smith, *A Manual of Methods for Baculovirus Vectors and Insect Cell Culture Procedures*, Texas Ag. Exper. Station Bulletin No. 1555 (1988), the entireties of which is herein incorporated by reference)

20 Methods for the expression of a nucleic acid sequence of interest in a fungal host cell are known in the art. The fungal host cell may, for example, be a yeast cell or a filamentous fungal cell. Methods for the expression of DNA sequences of interest in yeast cells are generally described in "Guide to yeast genetics and molecular biology", Guthrie and Fink, eds. *Methods in enzymology*, Academic Press, Inc. Vol 194 (1991) and *Gene expression technology*, Goeddel  
25 ed, *Methods in Enzymology*, Academic Press, Inc., Vol 185 (1991).

Mammalian cell lines available as hosts for expression are known in the art and include many immortalized cell lines available from the American Type Culture Collection (ATCC, Manassas, VA), such as HeLa cells, Chinese hamster ovary (CHO) cells, baby hamster kidney (BHK) cells and a number of other cell lines. Suitable promoters for mammalian cells are also  
30 known in the art and include, but are not limited to, viral promoters such as that from Simian



Virus 40 (SV40) (Fiers *et al.*, *Nature* 273:113 (1978), the entirety of which is herein incorporated by reference), Rous sarcoma virus (RSV), adenovirus (ADV) and bovine papilloma virus (BPV). Mammalian cells may also require terminator sequences and poly-A addition sequences. Enhancer sequences which increase expression may also be included and sequences which  
5 promote amplification of the gene may also be desirable (for example methotrexate resistance genes).

Vectors suitable for replication in mammalian cells are well known in the art, and may include viral replicons, or sequences which insure integration of the appropriate sequences encoding epitopes into the host genome. Plasmid vectors that greatly facilitate the construction of  
10 recombinant viruses have been described (*see*, for example, Mackett *et al.*, *J Virol.* 49:857 (1984); Chakrabarti *et al.*, *Mol. Cell. Biol.* 5:3403 (1985); Moss, In: *Gene Transfer Vectors For Mammalian Cells* (Miller and Calos, eds., Cold Spring Harbor Laboratory, N.Y., p. 10, (1987); all of which are herein incorporated by reference in their entirety).

The invention also includes plants and plant parts, such as seed, oil and meal derived  
15 from seed, and feed and food products processed from plants, which are enriched in tocopherols. Of particular interest is seed oil obtained from transgenic plants where the tocopherol level has been increased as compared to seed oil of a non-transgenic plant.

The harvested plant material may be subjected to additional processing to further enrich the tocopherol content. The skilled artisan will recognize that there are many such processes or  
20 methods for refining, bleaching and degumming oil. United States Patent Number 5,932,261, issued August 3, 1999, discloses on such process, for the production of a natural carotene rich refined and deodorised oil by subjecting the oil to a pressure of less than 0.060 mbar and to a temperature of less than 200.degree. C. Oil distilled by this process has reduced free fatty acids, yielding a refined, deodorised oil where Vitamin E contained in the feed oil is substantially  
25 retained in the processed oil. The teachings of this patent are incorporated herein by reference.

The invention now being generally described, it will be more readily understood by reference to the following examples which are included for purposes of illustration only and are not intended to limit the present invention.

## EXAMPLES

## Example 1: Identification of Prenyltransferase or tocopherol cyclase Sequences

5 PSI-BLAST (Altschul, *et al.* (1997) *Nuc Acid Res* 25:3389-3402) profiles were generated for both the straight chain and aromatic classes of prenyltransferases. To generate the straight chain profile, a prenyl-transferase from *Porphyra purpurea* (Genbank accession 1709766) was used as a query against the NCBI non-redundant protein database. The *E. coli* enzyme involved in the formation of ubiquinone, *ubiA* (genbank accession 1790473) was used as a starting  
10 sequence to generate the aromatic prenyltransferase profile. These profiles were used to search public and proprietary DNA and protein data bases. In *Arabidopsis* six putative prenyltransferases of the straight-chain class were identified, ATPT1, (SEQ ID NO:9), ATPT7 (SEQ ID NO:10), ATPT8 (SEQ ID NO:11), ATPT9 (SEQ ID NO:13), ATPT10 (SEQ ID NO:14), and ATPT11 (SEQ ID NO:15), and six were identified of the aromatic class, ATPT2  
15 (SEQ ID NO:1), ATPT3 (SEQ ID NO:3), ATPT4 (SEQ ID NO:5), ATPT5 (SEQ ID NO:7), ATPT6 (SEQ ID NO:8), and ATPT12 (SEQ ID NO:16). Additional prenyltransferase sequences from other plants related to the aromatic class of prenyltransferases, such as soy (SEQ ID NOs: 19-23, the deduced amino acid sequence of SEQ ID NO:23 is provided in SEQ ID NO:24) and maize (SEQ ID NOs:25-29, and 31) are also identified. The deduced amino acid sequence of  
20 ZMPT5 (SEQ ID NO:29) is provided in SEQ ID NO:30.

Searches are performed on a Silicon Graphics Unix computer using additional Bioaccelerator hardware and GenWeb software supplied by Compugen Ltd. This software and hardware enables the use of the Smith-Waterman algorithm in searching DNA and protein databases using profiles as queries. The program used to query protein databases is *profilesearch*.  
25 This is a search where the query is not a single sequence but a profile based on a multiple alignment of amino acid or nucleic acid sequences. The profile is used to query a sequence data set, i.e., a sequence database. The profile contains all the pertinent information for scoring each position in a sequence, in effect replacing the "scoring matrix" used for the standard query searches. The program used to query nucleotide databases with a protein profile is *tprofilesearch*.  
30 *Tprofilesearch* searches nucleic acid databases using an amino acid profile query. As the search is

running, sequences in the database are translated to amino acid sequences in six reading frames. The output file for tprofilesearch is identical to the output file for profilesearch except for an additional column that indicates the frame in which the best alignment occurred.

5 The Smith-Waterman algorithm, (Smith and Waterman (1981) *supra*), is used to search for similarities between one sequence from the query and a group of sequences contained in the database. E score values as well as other sequence information, such as conserved peptide sequences are used to identify related sequences.

10 To obtain the entire coding region corresponding to the *Arabidopsis* prenyltransferase sequences, synthetic oligo-nucleotide primers are designed to amplify the 5' and 3' ends of partial cDNA clones containing prenyltransferase sequences. Primers are designed according to the respective *Arabidopsis* prenyltransferase sequences and used in Rapid Amplification of cDNA Ends (RACE) reactions (Frohman *et al.* (1988) *Proc. Natl. Acad. Sci. USA* 85:8998-9002) using the Marathon cDNA amplification kit (Clontech Laboratories Inc, Palo Alto, CA).

15 Amino acid sequence alignments between ATPT2 (SEQ ID NO:2), ATPT3 (SEQ ID NO:4), ATPT4 (SEQ ID NO:6), ATPT8 (SEQ ID NO:12), and ATPT12 (SEQ ID NO:17) are performed using ClustalW (Figure 1), and the percent identity and similarities are provided in Table 1 below.

Table 1:

	ATPT2	ATPT3	ATPT4	ATPT8	ATPT12
ATPT2 % Identity		12	13	11	15
% similar		25	25	22	32
% Gap		17	20	20	9
ATPT3 % Identity			12	6	22
% similar			29	16	38
% Gap			20	24	14
ATPT4 % Identity				9	14
% similar				18	29
% Gap				26	19
ATPT8 % Identity					7

% similar	19
% Gap	20
ATPT12 % Identity	
% similar	
% Gap	

### Example 2: Preparation of Prenyl Transferase Expression Constructs

A plasmid containing the napin cassette derived from pCGN3223 (described in USPN 5,639,790, the entirety of which is incorporated herein by reference) was modified to make it more useful for cloning large DNA fragments containing multiple restriction sites, and to allow the cloning of multiple napin fusion genes into plant binary transformation vectors. An adapter comprised of the self annealed oligonucleotide of sequence CGCGATTAAATGGCGCGCCCTGCAGGCGGCCGCTGCAGGGCGCGCCATTAAAT (SEQ ID NO:40) was ligated into the cloning vector pBC SK+ (Stratagene) after digestion with the restriction endonuclease BssHII to construct vector pCGN7765. Plasmids pCGN3223 and pCGN7765 were digested with NotI and ligated together. The resultant vector, pCGN7770, contains the pCGN7765 backbone with the napin seed specific expression cassette from pCGN3223.

The cloning cassette, pCGN7787, essentially the same regulatory elements as pCGN7770, with the exception of the napin regulatory regions of pCGN7770 have been replaced with the double CAMV 35S promoter and the tml polyadenylation and transcriptional termination region.

A binary vector for plant transformation, pCGN5139, was constructed from pCGN1558 (McBride and Summerfelt, (1990) Plant Molecular Biology, 14:269-276). The polylinker of pCGN1558 was replaced as a HindIII/Asp718 fragment with a polylinker containing unique restriction endonuclease sites, AscI, PaeI, XbaI, SmaI, BamHI, and NotI. The Asp718 and HindIII restriction endonuclease sites are retained in pCGN5139.

A series of turbo binary vectors are constructed to allow for the rapid cloning of DNA sequences into binary vectors containing transcriptional initiation regions (promoters) and transcriptional termination regions.

The plasmid pCGN8618 was constructed by ligating oligonucleotides 5'-TCGAGGATCCGCGGCCGCAAGCTTCCTGCAGG-3' (SEQ ID NO:41) and 5'-TCGACCTGCAGGAAGCTTGCGGCCGCGGATCC-3' (SEQ ID NO:42) into Sall/XhoI-digested pCGN7770. A fragment containing the napin promoter, polylinker and napin 3' region was excised from pCGN8618 by digestion with Asp718I; the fragment was blunt-ended by filling in the 5' overhangs with Klenow fragment then ligated into pCGN5139 that had been digested with Asp718I and HindIII and blunt-ended by filling in the 5' overhangs with Klenow fragment. A plasmid containing the insert oriented so that the napin promoter was closest to the blunted Asp718I site of pCGN5139 and the napin 3' was closest to the blunted HindIII site was subjected to sequence analysis to confirm both the insert orientation and the integrity of cloning junctions. The resulting plasmid was designated pCGN8622.

The plasmid pCGN8619 was constructed by ligating oligonucleotides 5'-TCGACCTGCAGGAAGCTTGCGGCCGCGGATCC -3' (SEQ ID NO:43) and 5'-TCGAGGATCCGCGGCCGCAAGCTTCCTGCAGG-3' (SEQ ID NO:44) into Sall/XhoI-digested pCGN7770. A fragment containing the napin promoter, polylinker and napin 3' region was removed from pCGN8619 by digestion with Asp718I; the fragment was blunt-ended by filling in the 5' overhangs with Klenow fragment then ligated into pCGN5139 that had been digested with Asp718I and HindIII and blunt-ended by filling in the 5' overhangs with Klenow fragment. A plasmid containing the insert oriented so that the napin promoter was closest to the blunted Asp718I site of pCGN5139 and the napin 3' was closest to the blunted HindIII site was subjected to sequence analysis to confirm both the insert orientation and the integrity of cloning junctions. The resulting plasmid was designated pCGN8623.

The plasmid pCGN8620 was constructed by ligating oligonucleotides 5'-TCGAGGATCCGCGGCCGCAAGCTTCCTGCAGGAGCT -3' (SEQ ID NO:45) and 5'-CCTGCAGGAAGCTTGCGGCCGCGGATCC-3' (SEQ ID NO:46) into Sall/SacI-digested pCGN7787. A fragment containing the d35S promoter, polylinker and tml 3' region was removed from pCGN8620 by complete digestion with Asp718I and partial digestion with NotI. The fragment was blunt-ended by filling in the 5' overhangs with Klenow fragment then ligated into pCGN5139 that had been digested with Asp718I and HindIII and blunt-ended by filling in the 5' overhangs with Klenow fragment. A plasmid containing the insert oriented so that the



d35S promoter was closest to the blunted Asp718I site of pCGN5139 and the tml 3' was closest to the blunted HindIII site was subjected to sequence analysis to confirm both the insert orientation and the integrity of cloning junctions. The resulting plasmid was designated pCGN8624.

- 5       The plasmid pCGN8621 was constructed by ligating oligonucleotides 5'-TCGACCTGCAGGAAGCTTGCGGCCGCGGATCCAGCT -3' (SEQ ID NO:47) and 5'-GGATCCGCGGCCGCAAGCTTCCTGCAGG-3' (SEQ ID NO:48) into SalI/SacI-digested pCGN7787. A fragment containing the d35S promoter, polylinker and tml 3' region was removed from pCGN8621 by complete digestion with Asp718I and partial digestion with NotI.
- 10   The fragment was blunt-ended by filling in the 5' overhangs with Klenow fragment then ligated into pCGN5139 that had been digested with Asp718I and HindIII and blunt-ended by filling in the 5' overhangs with Klenow fragment. A plasmid containing the insert oriented so that the d35S promoter was closest to the blunted Asp718I site of pCGN5139 and the tml 3' was closest to the blunted HindIII site was subjected to sequence analysis to confirm both the insert
- 15   orientation and the integrity of cloning junctions. The resulting plasmid was designated pCGN8625.

- The plasmid construct pCGN8640 is a modification of pCGN8624 described above. A 938bp PstI fragment isolated from transposon Tn7 which encodes bacterial spectinomycin and streptomycin resistance (Fling et al. (1985), *Nucleic Acids Research* 13(19):7095-7106), a
- 20   determinant for E. coli and Agrobacterium selection, was blunt ended with Pfu polymerase. The blunt ended fragment was ligated into pCGN8624 that had been digested with SpeI and blunt ended with Pfu polymerase. The region containing the PstI fragment was sequenced to confirm both the insert orientation and the integrity of cloning junctions.

- The spectinomycin resistance marker was introduced into pCGN8622 and pCGN8623 as
- 25   follows. A 7.7 Kbp AvrII-SnaBI fragment from pCGN8640 was ligated to a 10.9 Kbp AvrII-SnaBI fragment from pCGN8623 or pCGN8622, described above. The resulting plasmids were pCGN8641 and pCGN8643, respectively.

      The plasmid pCGN8644 was constructed by ligating oligonucleotides 5'-GATCACCTGCAGGAAGCTTGCGGCCGCGGATCCAATGCA-3' (SEQ ID NO:49) and 5'-

TTGGATCCGCGGCCGCAAGCTTCCTGCAGGT-3' (SEQ ID NO:50) into BamHI-PstI digested pCGN8640.

Synthetic oligonucleotides were designed for use in Polymerase Chain Reactions (PCR) to amplify the coding sequences of ATPT2, ATPT3, ATPT4, ATPT8, and ATPT12 for the preparation of expression constructs and are provided in Table 2 below.

Table 2:

Name	Restriction Site	Sequence	SEQ ID NO:
ATPT2	5' NotI	GGATCCGCGGCCGCAACAATGGAGTC	51
		TCTGCTCTCTAGTTCT	
ATPT2	3' SseI	GGATCCTGCAGGTCACTTCAAAAAA	52
		GGTAACAGCAAGT	
ATPT3	5' NotI	GGATCCGCGGCCGCAACAATGGCGTT	53
		TTTTGGGCTCTCCCGTGTTT	
ATPT3	3' SseI	GGATCCTGCAGGTTATTGAAAACCTT	54
		CTTCCAAGTACAAC	
ATPT4	5' NotI	GGATCCGCGGCCGCAACAATGTGGCG	55
		AAGATCTGTTGTT	
ATPT4	3' SseI	GGATCCTGCAGGTCACTGGAGAGTAG	56
		AAGGAAGGAGCT	
ATPT8	5' NotI	GGATCCGCGGCCGCAACAATGGTACT	57
		TGCCGAGGTTCCAAAGCTTGCCTCT	
ATPT8	3' SseI	GGATCCTGCAGGTCACTTGTTTCTG	58
		GTGATGACTCTAT	
ATPT12	5' NotI	GGATCCGCGGCCGCAACAATGACTTC	59
		GATTCTCAACACT	
ATPT12	3' SseI	GGATCCTGCAGGTCACTGTTGCGAT	60
		GCTAATGCCGT	

The coding sequences of ATPT2, ATPT3, ATPT4, ATPT8, and ATPT12 were all amplified using the respective PCR primers shown in Table 2 above and cloned into the TopoTA vector (Invitrogen). Constructs containing the respective prenyltransferase sequences were digested with NotI and Sse8387I and cloned into the turbobinary vectors described above.

The sequence encoding ATPT2 prenyltransferase was cloned in the sense orientation into pCGN8640 to produce the plant transformation construct pCGN10800 (Figure 2). The ATPT2 sequence is under control of the 35S promoter.

The ATPT2 sequence was also cloned in the antisense orientation into the construct pCGN8641 to create pCGN10801 (Figure 3). This construct provides for the antisense expression of the ATPT2 sequence from the napin promoter.

5 The ATPT2 coding sequence was also cloned in the sense orientation into the vector pCGN8643 to create the plant transformation construct pCGN10822

The ATPT2 coding sequence was also cloned in the antisense orientation into the vector pCGN8644 to create the plant transformation construct pCGN10803 (Figure 4).

10 The ATPT4 coding sequence was cloned into the vector pCGN864 to create the plant transformation construct pCGN10806 (Figure 5). The ATPT2 coding sequence was cloned into the vector TopoTA™ vector from Invitrogen, to create the plant transformation construct pCGN10807 (Figure 6). The ATPT3 coding sequence was cloned into the TopoTA vector to create the plant transformation construct pCGN10808 (Figure 7). The ATPT3 coding sequence was cloned in the sense orientation into the vector pCGN8640 to create the plant transformation construct pCGN10809 (Figure 8). The ATPT3 coding sequence was cloned in the antisense orientation into the  
15 vector pCGN8641 to create the plant transformation construct pCGN10810 (Figure 9). The ATPT3 coding sequence was cloned into the vector pCGN8643 to create the plant transformation construct pCGN10811 (Figure 10). The ATPT3 coding sequence was cloned into the vector pCGN8644 to create the plant transformation construct pCGN10812 (Figure 11). The ATPT4 coding sequence was cloned into the vector pCGN8640 to create the plant transformation construct pCGN10813 (Figure  
20 12). The ATPT4 coding sequence was cloned into the vector pCGN8641 to create the plant transformation construct pCGN10814 (Figure 13). The ATPT4 coding sequence was cloned into the vector pCGN8643 to create the plant transformation construct pCGN10815 (Figure 14). The ATPT4 coding sequence was cloned in the antisense orientation into the vector pCGN8644 to create the plant transformation construct pCGN10816 (Figure 15). The ATPT8 coding sequence was cloned in  
25 the sense orientation into the vector pCGN8643 to create the plant transformation construct pCGN10819 (Figure 17). The ATPT12 coding sequence was cloned into the vector pCGN8640 to create the plant transformation construct pCGN10824 (Figure 18). The ATPT12 coding sequence was cloned into the vector pCGN8643 to create the plant transformation construct pCGN10825 (Figure 19). The ATPT8 coding sequence was cloned into the vector pCGN8640 to create the plant  
30 transformation construct pCGN10826 (Figure 20).

**Example 3: Plant Transformation with Prenyl Transferase Constructs**

Transgenic *Brassica* plants are obtained by *Agrobacterium*-mediated transformation as described by Radke *et al.* (*Theor. Appl. Genet.* (1988) 75:685-694; *Plant Cell Reports* (1992) 11:499-505). Transgenic *Arabidopsis thaliana* plants may be obtained by *Agrobacterium*-mediated transformation as described by Valverkens *et al.*, (*Proc. Nat. Acad. Sci.* (1988) 85:5536-5540), or as described by Bent *et al.* ((1994), *Science* 265:1856-1860), or Bechtold *et al.* ((1993), *C.R.Acad.Sci, Life Sciences* 316:1194-1199). Other plant species may be similarly transformed using related techniques.

Alternatively, microprojectile bombardment methods, such as described by Klein *et al.* (*Bio/Technology* 10:286-291) may also be used to obtain nuclear transformed plants.

**Example 4: Identification of Additional Prenyltransferases**

Additional BLAST searches were performed using the ATPT2 sequence, a sequence in the class of aromatic prenyltransferases. ESTs, and in some case, full-length coding regions, were identified in proprietary DNA libraries.

Soy full-length homologs to ATPT2 were identified by a combination of BLAST (using ATPT2 protein sequence) and 5' RACE. Two homologs resulted (SEQ ID NO:95 and SEQ ID NO:96). Translated amino acid sequences are provided by SEQ ID NO:97 and SEQ ID NO:98.

A rice est ATPT2 homolog is shown in SEQ ID NO:99 (obtained from BLAST using the wheat ATPT2 homolog).

Other homolog sequences were obtained using ATPT2 and PSI-BLAST, including est sequences from wheat (SEQ ID NO:100), leek (SEQ ID NOs:101 and 102), canola (SEQ ID NO:103), corn (SEQ ID NOs:104, 105 and 106), cotton (SEQ ID NO:107) and tomato (SEQ ID NO:108).

A PSI-Blast profile generated using the *E. coli* ubiA (genbank accession 1790473) sequence was used to analyze the *Synechocystis* genome. This analysis identified 5 open reading frames (ORFs) in the *Synechocystis* genome that were potentially prenyltransferases; slr0926 (annotated as ubiA (4-hydroxybenzoate-octaprenyltransferase, SEQ ID NO:32), slr1899

(annotated as *ctaB* (cytochrome c oxidase folding protein, SEQ ID NO:33), *slr0056* (annotated as *g4* (chlorophyll synthase 33 kd subunit, SEQ ID NO:34), *slr1518* (annotated as *menA* (menaquinone biosynthesis protein, SEQ ID NO:35), and *slr1736* (annotated as a hypothetical protein of unknown function (SEQ ID NO:36).

5

#### 4A. *Synechocystis* Knock-outs

To determine the functionality of these ORFs and their involvement, if any, in the biosynthesis of tocopherols, knockouts constructs were made to disrupt the ORF identified in *Synechocystis*.

10

Synthetic oligos were designed to amplify regions from the 5' (5'-TAATGTGTACATTGTCGGCCTC (17365') (SEQ ID NO:61) and 5'-GCAATGTAACATCAGAGATTTTGAGACACAACGTGGCTTTCCACAATTCCCCGCACCGTC (1736kanpr1)) (SEQ ID NO:62) and 3' (5'-AGGCTAATAAGCACAAATGGGA (17363') (SEQ ID NO:63) and 5'-GGTATGAGTCAGCAACACCTTCTTCACGAGGCAGACCTCAGCGGAATTGGTTTAGGTTATCCC (1736kanpr2)) (SEQ ID NO:64) ends of the *slr1736* ORF.

15

The 1736kanpr1 and 1736kanpr2 oligos contained 20 bp of homology to the *slr1736* ORF with an additional 40 bp of sequence homology to the ends of the kanamycin resistance cassette.

Separate PCR steps were completed with these oligos and the products were gel purified and combined with the kanamycin resistance gene from puc4K (Pharmacia) that had been digested

20

with *HincII* and gel purified away from the vector backbone. The combined fragments were allowed to assemble without oligos under the following conditions: 94°C for 1 min, 55°C for 1 min, 72°C for 1 min plus 5 seconds per cycle for 40 cycles using pfu polymerase in 100ul reaction volume (Zhao, H and Arnold (1997) *Nucleic Acids Res.* 25(6):1307-1308). One

25

microliter or five microliters of this assembly reaction was then amplified using 5' and 3' oligos nested within the ends of the ORF fragment, so that the resulting product contained 100-200 bp of the 5' end of the *Synechocystis* gene to be knocked out, the kanamycin resistance cassette, and 100-200 bp of the 3' end of the gene to be knocked out. This PCR product was then cloned into the vector pGemT easy (Promega) to create the construct pMON21681 and used for *Synechocystis* transformation.



Primers were also synthesized for the preparation of *Synechocystis* knockout constructs for the other sequences using the same method as described above, with the following primers.

The *ubiA* 5' sequence was amplified using the primers 5'- GGATCCATGGTT  
GCCCCAACCCTATC (SEQ ID NO:65) and 5'- GCAATGTAAACATCAGAGA

5 TTTTGAGACACAACG TGGCTTTGGGTAAAGCAACAATGACCGGC (SEQ ID NO:66).

The 3' region was amplified using the synthetic oligonucleotide primers 5'-

GAATTCTCAAAGCCAGCCCAGTAAC (SEQ ID NO:67) and 5'-GGTATGAGTC  
AGCAACACCTTCTTCACGAGGCAGACCTCAGCGGGTGCGAAAAGGGTTTTCCC (SEQ  
ID NO:68). The amplification products were combined with the kanamycin resistance gene from

10 puc4K (Pharmacia) that had been digested with *HincII* and gel purified away from the vector  
backbone. The annealed fragment was amplified using 5' and 3' oligos nested within the ends of  
the ORF fragment (5'- CCAGTGGTTTAGGCTGTGTGGTC (SEQ ID NO:69) and 5'-  
CTGAGTTGGATGTATTGGATC (SEQ ID NO:70)), so that the resulting product contained  
100-200 bp of the 5' end of the *Synechocystis* gene to be knocked out, the kanamycin resistance  
15 cassette, and 100-200 bp of the 3' end of the gene to be knocked out. This PCR product was then  
cloned into the vector pGemT easy (Promega) to create the construct pMON21682 and used for  
*Synechocystis* transformation.

Primers were also synthesized for the preparation of *Synechocystis* knockout constructs for the other sequences using the same method as described above, with the following primers.

20 The *sl11899* 5' sequence was amplified using the primers 5'- GGATCCATGGTTACTT  
CGACAAAATCC (SEQ ID NO:71) and 5'- GCAATGTAAACATCAGAG  
ATTTTGAGACACAACGTGGCTTTGCTAGGCAACCGCTTAGTAC (SEQ ID NO:72). The  
3' region was amplified using the synthetic oligonucleotide primers 5'-

GAATTCTTAACCCAACAGTAAAGTTCCC (SEQ ID NO:73) and 5'- GGTATGAGTCAGC  
25 AACACCTTCTTCACGAGGCAGACCTCAGCGCCGGCATTGTCTTTTACATG (SEQ ID  
NO:74). The amplification products were combined with the kanamycin resistance gene from  
puc4K (Pharmacia) that had been digested with *HincII* and gel purified away from the vector  
backbone. The annealed fragment was amplified using 5' and 3' oligos nested within the ends of  
the ORF fragment (5'- GGAACCCTTGCAGCCGCTTC (SEQ ID NO:75)

and 5'-GTATGCCCAACTGGTGCAGAGG (SEQ ID NO:76)), so that the resulting product contained 100-200 bp of the 5' end of the *Synechocystis* gene to be knocked out, the kanamycin resistance cassette, and 100-200 bp of the 3' end of the gene to be knocked out. This PCR product was then cloned into the vector pGemT easy (Promega) to create the construct pMON21679 and used for *Synechocystis* transformation.

Primers were also synthesized for the preparation of *Synechocystis* knockout constructs for the other sequences using the same method as described above, with the following primers.

The slr0056 5' sequence was amplified using the primers 5'-

GGATCCATGTCTGACACACAAAATACCG (SEQ ID NO:77) and 5'-

10 GCAATGTAACATCAGAGATTTTGAGACACAACGTGGCTTTCGCCAATACCAGCCACC AACAG (SEQ ID NO:78). The 3' region was amplified using the synthetic oligonucleotide

primers 5'-GAATTCTCAAATCCCCGCATGGCCTAG (SEQ ID NO:79) and 5'-

GGTATGAGTCAGCAACACCTTCTTCACGAGGCAGACCTCAGCGGCCTACGGCTTGGA CGTGTGGG (SEQ ID NO:80). The amplification products were combined with the kanamycin

15 resistance gene from puc4K (Pharmacia) that had been digested with *HincII* and gel purified away from the vector backbone. The annealed fragment was amplified using 5' and 3' oligos nested within the ends of the ORF fragment (5'-CACTTGGATTCCCCTGATCTG (SEQ ID NO:81) and 5'-GCAATACCCGCTTGGAACG (SEQ ID NO:82)), so that the resulting product contained 100-200 bp of the 5' end of the *Synechocystis* gene to be knocked out, the  
20 kanamycin resistance cassette, and 100-200 bp of the 3' end of the gene to be knocked out. This PCR product was then cloned into the vector pGemT easy (Promega) to create the construct pMON21677 and used for *Synechocystis* transformation.

Primers were also synthesized for the preparation of *Synechocystis* knockout constructs for the other sequences using the same method as described above, with the following primers.

25 The slr1518 5' sequence was amplified using the primers 5'-GGATCCATGACCGAAT CTTGCCCCCTAGC (SEQ ID NO:83) and 5'-GCAATGTAACATCAGAGATTTTGA GACACAACGTGGC TTTCAATCCTAGGTAGCCGAGGCG (SEQ ID NO:84). The 3' region was amplified using the synthetic oligonucleotide primers 5'-GAATTCTTAGCCCAGGCC AGCCCAGCC (SEQ ID NO:85) and 5'-GGTATGAGTCAGCAACACCTTCTTCACGA  
30 GGCAGACCTCAGCGGGGAATTGATTTGTTTAATTACC (SEQ ID NO:86). The

amplification products were combined with the kanamycin resistance gene from puc4K (Pharmacia) that had been digested with *HincII* and gel purified away from the vector backbone. The annealed fragment was amplified using 5' and 3' oligos nested within the ends of the ORF fragment (5'- GCGATCGCCATTATCGCTTGG (SEQ ID NO:87) and 5'-  
5 GCAGACTGGCAATTATCAGTAACG (SEQ ID NO:88)), so that the resulting product contained 100-200 bp of the 5' end of the *Synechocystis* gene to be knocked out, the kanamycin resistance cassette, and 100-200 bp of the 3' end of the gene to be knocked out. This PCR product was then cloned into the vector pGcmT easy (Promega) to create the construct pMON21680 and used for *Synechocystis* transformation.

10

#### 4B. Transformation of *Synechocystis*

Cells of *Synechocystis* 6803 were grown to a density of approximately  $2 \times 10^8$  cells per ml and harvested by centrifugation. The cell pellet was re-suspended in fresh BG-11 medium (ATCC Medium 616) at a density of  $1 \times 10^9$  cells per ml and used immediately for transformation.

15 One-hundred microliters of these cells were mixed with 5 ul of mini prep DNA and incubated with light at 30C for 4 hours. This mixture was then plated onto nylon filters resting on BG-11 agar supplemented with TES pH8 and allowed to grow for 12-18 hours. The filters were then transferred to BG-11 agar + TES + 5ug/ml kanamycin and allowed to grow until colonies appeared within 7-10 days (Packer and Glazer, 1988). Colonies were then picked into BG-11  
20 liquid media containing 5 ug/ml kanamycin and allowed to grow for 5 days. These cells were then transferred to Bg-11 media containing 10ug/ml kanamycin and allowed to grow for 5 days and then transferred to Bg-11 + kanamycin at 25ug/ml and allowed to grow for 5 days. Cells were then harvested for PCR analysis to determine the presence of a disrupted ORF and also for HPLC analysis to determine if the disruption had any effect on tocopherol levels.

25 PCR analysis of the *Synechocystis* isolates for slr1736 and sl11899 showed complete segregation of the mutant genome, meaning no copies of the wild type genome could be detected in these strains. This suggests that function of the native gene is not essential for cell function. HPLC analysis of these same isolates showed that the sl11899 strain had no detectable reduction in tocopherol levels. However, the strain carrying the knockout for slr1736 produced no  
30 detectable levels of tocopherol.

The amino acid sequences for the *Synechocystis* knockouts are compared using ClustalW, and are provided in Table 3 below. Provided are the percent identities, percent similarity, and the percent gap. The alignment of the sequences is provided in Figure 21.

5 **Table 3:**

	Slr1736	slr0926	sll1899	slr0056	slr1518
slr1736 %identity		14	12	18	11
%similar		29	30	34	26
%gap		8	7	10	5
slr0926 %identity			20	19	14
%similar			39	32	28
%gap			7	9	4
sll1899 %identity				17	13
%similar				29	29
%gap				12	9
slr0056 %identity					15
%similar					31
%gap					8
slr1518 %identity					
%similar					
%gap					

Amino acid sequence comparisons are performed using various *Arabidopsis* prenyltransferase sequences and the *Synechocystis* sequences. The comparisons are presented in Table 4 below. Provided are the percent identities, percent similarity, and the percent gap. The alignment of the sequences is provided in Figure 22.

10 **Table 4:**

	ATPT2	slr1736	ATPT3	slr0926	ATPT4	sll1899	ATPT12	slr0056	ATPT8	slr1518
ATPT2		29	9	9	8	8	12	9	7	9

	46	23	21	20	20	28	23	21	20
	27	13	28	23	29	11	24	25	24
slr1736		9	13	8	12	13	15	8	10
		19	28	19	28	26	33	21	26
		34	12	34	15	26	10	12	10
ATPT3			23	11	14	13	10	5	11
			36	26	26	26	21	14	22
			29	21	31	16	30	30	30
				12	20	17	20	11	14
slr0926				24	37	28	33	24	29
				33	12	25	10	11	9
					18	11	8	6	7
ATPT4					33	23	18	16	19
					28	19	32	32	33
						13	17	10	12
slr1899						24	30	23	26
						27	13	10	11
							52	8	11
ATPT1							66	19	26
2									
							18	25	23
								9	13
slr0056								23	32
								10	8
									7
ATPT8									23
									7
slr1518									





#### 4C. Phytyl Prenyltransferase Enzyme Assays

[<sup>3</sup>H] Homogentisic acid in 0.1% H<sub>3</sub>PO<sub>4</sub> (specific radioactivity 40 Ci/mmol). Phytyl  
5 pyrophosphate was synthesized as described by Joo, *et al.* (1973) *Can J. Biochem.* 51:1527. 2-methyl-6-phytylquinol and 2,3-dimethyl-5-phytylquinol were synthesized as described by Soll, *et al.* (1980) *Phytochemistry* 19:215. Homogentisic acid,  $\alpha$ ,  $\beta$ ,  $\delta$ , and  $\gamma$ -tocopherol, and tocol, were purchased commercially.

The wild-type strain of *Synechocystis* sp. PCC 6803 was grown in BG11 medium with  
10 bubbling air at 30°C under 50  $\mu\text{E}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$  fluorescent light, and 70% relative humidity. The growth medium of slr1736 knock-out (potential PPT) strain of this organism was supplemented with 25  $\mu\text{g mL}^{-1}$  kanamycin. Cells were collected from 0.25 to 1 liter culture by centrifugation at 5000 *g* for 10 min and stored at -80°C.

Total membranes were isolated according to Zak's procedures with some modifications (Zak,  
15 *et al.* (1999) *Eur J. Biochem* 261:311). Cells were broken on a French press. Before the French press treatment, the cells were incubated for 1 hour with lysozyme (0.5%, w/v) at 30 °C in a medium containing 7 mM EDTA, 5 mM NaCl and 10 mM Hepes-NaOH, pH 7.4. The spheroplasts were collected by centrifugation at 5000 *g* for 10 min and resuspended at 0.1 - 0.5 mg chlorophyll·mL<sup>-1</sup> in 20 mM potassium phosphate buffer, pH 7.8. Proper amount of protease  
20 inhibitor cocktail and DNAase I from Boehringer Mannheim were added to the solution. French press treatments were performed two to three times at 100 MPa. After breakage, the cell suspension was centrifuged for 10 min at 5000*g* to pellet unbroken cells, and this was followed by centrifugation at 100 000 *g* for 1 hour to collect total membranes. The final pellet was resuspended in a buffer containing 50 mM Tris-HCL and 4 mM MgCl<sub>2</sub>.

25 Chloroplast pellets were isolated from 250 g of spinach leaves obtained from local markets. Devined leaf sections were cut into grinding buffer (2 l /250 g leaves) containing 2 mM EDTA, 1 mM MgCl<sub>2</sub>, 1 mM MnCl<sub>2</sub>, 0.33 M sorbitol, 0.1% ascorbic acid, and 50 mM Hepes at pH 7.5. The leaves were homogenized for 3 sec three times in a 1-L blender, and filtered through 4 layers of miracloth. The supernatant was then centrifuged at 5000*g* for 6 min. The chloroplast pellets were

resuspended in small amount of grinding buffer (Douce, *et al* Methods in Chloroplast Molecular Biology, 239 (1982))

Chloroplasts in pellets can be broken in three ways. Chloroplast pellets were first aliquoted in 1 mg of chlorophyll per tube, centrifuged at 6000 rpm for 2 min in microcentrifuge, and grinding buffer was removed. Two hundred microliters of Triton X-100 buffer (0.1% Triton X-100, 50 mM Tris-HCl pH 7.6 and 4 mM MgCl<sub>2</sub>) or swelling buffer (10 mM Tris pH 7.6 and 4 mM MgCl<sub>2</sub>) was added to each tube and incubated for ½ hour at 4°C. Then the broken chloroplast pellets were used for the assay immediately. In addition, broken chloroplasts can also be obtained by freezing in liquid nitrogen and stored at -80°C for ½ hour, then used for the assay.

In some cases chloroplast pellets were further purified with 40%/ 80% percoll gradient to obtain intact chloroplasts. The intact chloroplasts were broken with swelling buffer, then either used for assay or further purified for envelope membranes with 20.5%/ 31.8% sucrose density gradient (Sol, *et al* (1980) *supra*). The membrane fractions were centrifuged at 100 000g for 40 min and resuspended in 50 mM Tris-HCl pH 7.6, 4 mM MgCl<sub>2</sub>.

Various amounts of [<sup>3</sup>H]HGA, 40 to 60 µM unlabelled HGA with specific activity in the range of 0.16 to 4 Ci/mmol were mixed with a proper amount of 1M Tris-NaOH pH 10 to adjust pH to 7.6. HGA was reduced for 4 min with a trace amount of solid NaBH<sub>4</sub>. In addition to HGA, standard incubation mixture (final vol 1 mL) contained 50 mM Tris-HCl, pH 7.6, 3-5 mM MgCl<sub>2</sub>, and 100 µM phytyl pyrophosphate. The reaction was initiated by addition of *Synechocystis* total membranes, spinach chloroplast pellets, spinach broken chloroplasts, or spinach envelope membranes. The enzyme reaction was carried out for 2 hour at 23°C or 30°C in the dark or light. The reaction is stopped by freezing with liquid nitrogen, and stored at -80°C or directly by extraction.

A constant amount of tocol was added to each assay mixture and reaction products were extracted with a 2 mL mixture of chloroform/methanol (1:2, v/v) to give a monophasic solution. NaCl solution (2 mL; 0.9%) was added with vigorous shaking. This extraction procedure was repeated three times. The organic layer containing the prenylquinones was filtered through a 20 µm filter, evaporated under N<sub>2</sub> and then resuspended in 100 µL of ethanol.

The samples were mainly analyzed by Normal-Phase HPLC method (Isocratic 90% Hexane and 10% Methyl-t-butyl ether), and use a Zorbax silica column, 4.6 x 250 mm. The samples were

also analyzed by Reversed-Phase HPLC method (Isocratic 0.1%  $\text{H}_3\text{PO}_4$  in MeOH), and use a Vydac 2011IS54 C18 column; 4.6 x 250 mm coupled with an All-tech C18 guard column. The amount of products were calculated based on the substrate specific radioactivity, and adjusted according to the % recovery based on the amount of internal standard.

- 5        The amount of chlorophyll was determined as described in Arnon (1949) *Plant Physiol.* 24:1. Amount of protein was determined by the Bradford method using gamma globulin as a standard (Bradford, (1976) *Anal. Biochem.* 72:248)

10        Results of the assay demonstrate that 2-Methyl-6-Phytylplastoquinone is not produced in the *Synechocystis* slr1736 knockout preparations. The results of the phytyl prenyltransferase enzyme activity assay for the slr1736 knock out are presented in Figure 23.

#### 4D. Complementation of the slr1736 knockout with ATPT2

15        In order to determine whether ATPT2 could complement the knockout of slr1736 in *Synechocystis* 6803, a plasmid was constructed to express the ATPT2 sequence from the TAC promoter. A vector, plasmid psl1211, was obtained from the lab of Dr. Himadri Pakrasi of Washington University, and is based on the plasmid RSF1010 which is a broad host range plasmid (Ng W.-O., Zentella R., Wang, Y., Taylor J-S. A., Pakrasi, H.B. 2000. *phrA*, the major photoreactivating factor in the cyanobacterium *Synechocystis* sp. strain PCC 6803 codes for a cyclobutane pyrimidine dimer specific DNA photolyase. *Arch. Microbiol.* (in press)). The

20        ATPT2 gene was isolated from the vector pCGN10817 by PCR using the following primers. ATPT2nco.pr 5'-CCATGGATTTCGAGTAAAGTTGTCGC (SEQ ID NO:89); ATPT2ri.pr- 5'-GAATTCACCTCAAAAAAGGTAACAG (SEQ ID NO:90). These primers will remove approximately 112 BP from the 5' end of the ATPT2 sequence, which is thought to be the chloroplast transit peptide. These primers will also add an NcoI site at the 5' end and an EcoRI

25        site at the 3' end which can be used for sub-cloning into subsequent vectors. The PCR product from using these primers and pCGN10817 was ligated into pGEM T easy and the resulting vector pMON21689 was confirmed by sequencing using the m13forward and m13reverse primers. The NcoI/EcoRI fragment from pMON21689 was then ligated with the EagI/EcoRI and EagI/NcoI fragments from psl1211 resulting in pMON21690. The plasmid pMON21690

30        was introduced into the slr1736 *Synechocystis* 6803 KO strain via conjugation. Cells of sl906 (a

helper strain) and DH10B cells containing pMON21690 were grown to log phase (O.D. 600 = 0.4) and 1 ml was harvested by centrifugation. The cell pellets were washed twice with a sterile BG-11 solution and resuspended in 200 ul of BG-11. The following was mixed in a sterile eppendorf tube: 50 ul SL906, 50 ul DH10B cells containing pMON21690, and 100 ul of a fresh culture of the slr1736 *Synechocystis* 6803 KO strain (O.D. 730 = 0.2-0.4). The cell mixture was immediately transferred to a nitrocellulose filter resting on BG-11 and incubated for 24 hours at 30C and 2500 LUX(50 ue) of light. The filter was then transferred to BG-11 supplemented with 10ug/ml Gentamycin and incubated as above for ~5 days. When colonies appeared, they were picked and grown up in liquid BG-11 + Gentamycin 10 ug/ml. (Elhai, J. and Wolk, P. 1988.

10 Conjugal transfer of DNA to Cyanobacteria. *Methods in Enzymology* 167, 747-54) The liquid cultures were then assayed for tocopherols by harvesting 1ml of culture by centrifugation, extracting with ethanol/pyrogallol, and HPLC separation. The slr1736 *Synechocystis* 6803 KO strain, did not contain any detectable tocopherols, while the slr1736 *Synechocystis* 6803 KO strain transformed with pmon21690 contained detectable alpha tocopherol. A *Synechocystis* 15 6803 strain transformed with psl1211(vector control) produced alpha tocopherol as well.

#### 4E: Additional Evidence of Prenyltransferase Activity

To test the hypothesis that slr1736 or ATPT2 are sufficient as single genes to obtain phytyl prenyltransferase activity, both genes were expressed in SF9 cells and in yeast. When either slr1736 or ATPT2 were expressed in insect cells (Table 5) or in yeast, phytyl prenyltransferase activity was detectable in membrane preparations, whereas membrane preparations of the yeast vector control, or membrane preparations of insect cells did not exhibit phytyl prenyltransferase activity.

Table 5: Phytyl prenyltransferase activity

Enzyme source	Enzyme activity [pmol/mg x h]
slr1736 expressed in SF9 cells	20
ATPT2 expressed in SF9 cells	6
SF9 cell control	< 0.05

<i>Synechocystis</i> 6803	0.25
Spinach chloroplasts	0.20

### Example 5: Transgenic Plant Analysis

#### 5A. *Arabidopsis*

5 *Arabidopsis* plants transformed with constructs for the sense or antisense expression of the ATPT proteins were analyzed by High Pressure Liquid Chromatography (HPLC) for altered levels of total tocopherols, as well as altered levels of specific tocopherols (alpha, beta, gamma, and delta tocopherol).

10 Extracts of leaves and seeds were prepared for HPLC as follows. For seed extracts, 10 mg of seed was added to 1 g of microbeads (Biospec) in a sterile microfuge tube to which 500 ul 1% pyrogallol (Sigma Chem)/ethanol was added. The mixture was shaken for 3 minutes in a mini Beadbeater (Biospec) on "fast" speed. The extract was filtered through a 0.2 um filter into an autosampler tube. The filtered extracts were then used in HPLC analysis described below.

15 Leaf extracts were prepared by mixing 30-50 mg of leaf tissue with 1 g microbeads and freezing in liquid nitrogen until extraction. For extraction, 500 ul 1% pyrogallol in ethanol was added to the leaf/bead mixture and shaken for 1 minute on a Beadbeater (Biospec) on "fast" speed. The resulting mixture was centrifuged for 4 minutes at 14,000 rpm and filtered as described above prior to HPLC analysis.

20 HPLC was performed on a Zorbax silica HPLC column (4.6 mm X 250 mm) with a fluorescent detection, an excitation at 290 nm, an emission at 336 nm, and bandpass and slits. Solvent A was hexane and solvent B was methyl-t-butyl ether. The injection volume was 20 ul, the flow rate was 1.5 ml/min, the run time was 12 min (40°C) using the gradient (Table 6):

Table 6:

25	<u>Time</u>	<u>Solvent A</u>	<u>Solvent B</u>
	0 min.	90%	10%
	10 min.	90%	10%
	11 min.	25%	75%
	12 min.	90%	10%



Tocopherol standards in 1% pyrogallol/ ethanol were also run for comparison (alpha tocopherol, gamma tocopherol, beta tocopherol, delta tocopherol, and tocopherol (tocol) (all from Matreya).

5 Standard curves for alpha, beta, delta, and gamma tocopherol were calculated using Chemstation software. The absolute amount of component x is: Absolute amount of x =  $\text{Response}_x \times \text{RF}_x \times \text{dilution factor}$  where  $\text{Response}_x$  is the area of peak x,  $\text{RF}_x$  is the response factor for component x ( $\text{Amount}_x / \text{Response}_x$ ) and the dilution factor is 500 ul. The ng/mg tissue is found by: total ng component/mg plant tissue.

10 Results of the HPLC analysis of seed extracts of transgenic *Arabidopsis* lines containing pMON10822 for the expression of ATPT2 from the napin promoter are provided in Figure 24.

HPLC analysis results of segregating T2 *Arabidopsis* seed tissue expressing the ATPT2 sequence from the napin promoter (pCGN10822) demonstrates an increased level of tocopherols in the seed. Total tocopherol levels are increased as much as 50% over the total tocopherol  
15 levels of non-transformed (wild-type) *Arabidopsis* plants (Figure 25). Homozygous progeny from the top 3 lines (T3 seed) have up to a two-fold (100%) increase in total tocopherol levels over control *Arabidopsis* seed ( Figure 26.)

Furthermore, increases of particular tocopherols are also increased in transgenic *Arabidopsis* plants expressing the ATPT2 nucleic acid sequence from the napin promoter.

20 Levels of delta tocopherol in these lines are increased greater than 3 fold over the delta tocopherol levels obtained from the seeds of wild type *Arabidopsis* lines. Levels of gamma tocopherol in transgenic *Arabidopsis* lines expressing the ATPT2 nucleic acid sequence are increased as much as about 60% over the levels obtained in the seeds of non-transgenic control lines. Furthermore, levels of alpha tocopherol are increased as much as 3 fold over those  
25 obtained from non-transgenic control lines.

Results of the HPLC analysis of seed extracts of transgenic *Arabidopsis* lines containing pCGN10803 for the expression of ATPT2 from the enhanced 35S promoter (antisense orientation ) are provided in Figure 25. Two lines were identified that have reduced total tocopherols, up to a ten-fold decrease observed in T3 seed compared to control *Arabidopsis*  
30 (Figure 27.)

## 5B. Canola

Brassica napus, variety SP30021, was transformed with pCGN10822 (napin-A1PT2-napin 3', sense orientation) using *Agrobacterium tumefaciens*-mediated transformation. Flowers of the R0 plants were tagged upon pollination and developing seed was collected at 35 and 45 days after pollination (DAP).

Developing seed was assayed for tocopherol levels, as described above for *Arabidopsis*. Line 10822-1 shows a 20% increase of total tocopherols, compared to the wild-type control, at 45 DAP. Figure 28 shows total tocopherol levels measured in developing canola seed.

## Example 6: Sequences to Tocopherol Cyclase

### 6A. Preparation of the *slr1737* Knockout

The *Synechocystis* sp. 6803 *slr1737* knockout was constructed by the following method. The GPS™-1 Genome Priming System (New England Biolabs) was used to insert, by a Tn7 Transposase system, a Kanamycin resistance cassette into *slr1737*. A plasmid from a *Synechocystis* genomic library clone containing 652 base pairs of the targeted orf (*Synechocystis* genome base pairs 1324051 – 1324703; the predicted orf base pairs 1323672 – 1324763, as annotated by Cyanobase) was used as target DNA. The reaction was performed according to the manufacturers protocol. The reaction mixture was then transformed into *E. coli* DH10B electrocompetant cells and plated. Colonies from this transformation were then screened for transposon insertions into the target sequence by amplifying with M13 Forward and Reverse Universal primers, yielding a product of 652 base pairs plus ~1700 base pairs, the size of the transposon kanamycin cassette, for a total fragment size of ~2300 base pairs. After this determination, it was then necessary to determine the approximate location of the insertion within the targeted orf, as 100 base pairs of orf sequence was estimated as necessary for efficient homologous recombination in *Synechocystis*. This was accomplished through amplification reactions using either of the primers to the ends of the transposon, Primer S (5' end) or N (3' end), in combination with either a M13 Forward or Reverse primer. That is, four different primer combinations were used to map each potential knockout construct: Primer S – M13 Forward, Primer S – M13 Reverse, Primer N – M13 Forward, Primer N – M13 Reverse. The construct

used to transform *Synechocystis* and knockout *slr1737* was determined to consist of a approximately 150 base pairs of *slr1737* sequence on the 5' side of the transposon insertion and approximately 500 base pairs on the 3' side, with the transcription of the orf and kanamycin cassette in the same direction. The nucleic acid sequence of *slr1737* is provided in SEQ ID NO:38 the deduced amino acid sequence is provided in SEQ ID NO:39.

Cells of *Synechocystis* 6803 were grown to a density of  $\sim 2 \times 10^8$  cells per ml and harvested by centrifugation. The cell pellet was re-suspended in fresh BG-11 medium at a density of  $1 \times 10^9$  cells per ml and used immediately for transformation. 100 ul of these cells were mixed with 5 ul of mini prep DNA and incubated with light at 30C for 4 hours. This mixture was then plated onto nylon filters resting on BG-11 agar supplemented with TES pH8 and allowed to grow for 12-18 hours. The filters were then transferred to BG-11 agar + TES + 5ug/ml kanamycin and allowed to grow until colonies appeared within 7-10 days (Packer and Glazer, 1988). Colonies were then picked into BG-11 liquid media containing 5 ug/ml kanamycin and allowed to grow for 5 days. These cells were then transferred to Bg-11 media containing 10ug/ml kanamycin and allowed to grow for 5 days and then transferred to Bg-11 + kanamycin at 25ug/ml and allowed to grow for 5 days. Cells were then harvested for PCR analysis to determine the presence of a disrupted ORF and also for HPLC analysis to determine if the disruption had any effect on tocopherol levels.

PCR analysis of the *Synechocystis* isolates, using primers to the ends of the *slr1737* orf, showed complete segregation of the mutant genome, meaning no copies of the wild type genome could be detected in these strains. This suggests that function of the native gene is not essential for cell function. HPLC analysis of the strain carrying the knockout for *slr1737* produced no detectable levels of tocopherol.

#### 6B. The relation of *slr1737* and *slr1736*

The *slr1737* gene occurs in *Synechocystis* downstream and in the same orientation as *slr1736*, the phytol prenyltransferase. In bacteria this proximity often indicates an operon structure and therefore an expression pattern that is linked in all genes belonging to this operon. Occasionally such operons contain several genes that are required to constitute one enzyme. To confirm that *slr1737* is not required for phytol prenyltransferase activity, phytol prenyltransferase

was measured in extracts from the *Synechocystis* slr1737 knockout mutant. Figure 29 shows that extracts from the *Synechocystis* slr1737 knockout mutant still contain phytyl prenyltransferase activity. The molecular organization of genes in *Synechocystis* 6803 is shown in A. Figures B and C show HPLC traces (normal phase HPLC) of reaction products obtained with membrane preparations from *Synechocystis* wild type and slr1737<sup>-</sup> membrane preparations, respectively.

The fact that slr1737 is not required for the PPT activity provides additional data that ATPT2 and slr1736 encode phytyl prenyltransferases.

#### 6C *Synechocystis* Knockouts

*Synechocystis* 6803 wild type and *Synechocystis* slr1737 knockout mutant were grown photoautotrophically. Cells from a 20 ml culture of the late logarithmic growth phase were harvested and extracted with ethanol. Extracts were separated by isocratic normal-phase HPLC using a Hexane/Methyl-t-butyl ether (95/5) and a Zorbax silica column, 4.6 x 250 mm. Tocopherols and tocopherol intermediates were detected by fluorescence (excitement 290 nm, emission 336 nm) (Figure 30).

Extracts of *Synechocystis* 6803 contained a clear signal of alpha-tocopherol. 2,3-Dimethyl-5-phytylplastoquinol was below the limit of detection in extracts from the *Synechocystis* wild type (C). In contrast, extracts from the *Synechocystis* slr1737 knockout mutant did not contain alpha-tocopherol, but contained 2,3-dimethyl-5-phytylplastoquinol (D), indicating that the interruption of slr1737 has resulted in a block of the 2,3-dimethyl-5-phytylplastoquinol cyclase reaction.

Chromatograms of standard compounds alpha, beta, gamma, delta-tocopherol and 2,3-dimethyl-5-phytylplastoquinol are shown in A and B. Chromatograms of extracts from *Synechocystis* wild type and the *Synechocystis* slr1737 knockout mutant are shown in C and D, respectively. Abbreviations: 2,3-DMPQ, 2,3-dimethyl-5-phytylplastoquinol.

#### 6D. Incubation with Lysozyme treated *Synechocystis*

*Synechocystis* 6803 wild type and slr1737 knockout mutant cells from the late logarithmic growth phase (approximately 1g wet cells per experiment in a total volume of 3 ml) were treated with Lysozyme and subsequently incubated with S-adenosylmethionine, and

phytylpyrophosphate, plus radiolabelled homogentisic acid. After 17h incubation in the dark at room temperature the samples were extracted with 6 ml chloroform / methanol (1/2 v/v). Phase separation was obtained by the addition of 6 ml 0.9% NaCl solution. This procedure was repeated three times. Under these conditions 2,3-dimethyl-5-phytylplastoquinol is oxidized to  
5 form 2,3-dimethyl-5-phytylplastoquinone.

The extracts were analyzed by normal phase and reverse phase HPLC. Using extracts from wild type *Synechocystis* cells radiolabelled gamma-tocopherol and traces of radiolabelled 2,3-dimethyl-5-phytylplastoquinone were detected. When extracts from the slr1737 knockout mutant were analyzed, only radiolabelled 2,3-dimethyl-5-phytylplastoquinone was detectable.

- 10 The amount of 2,3-dimethyl-5-phytylplastoquinone was significantly increased compared to wild type extracts. Heat treated samples of the wild type and the slr1737 knockout mutant did not produce radiolabelled 2,3-dimethyl-5-phytylplastoquinone, nor radiolabelled tocopherols. These results further support the role of the slr1737 expression product in the cyclization of 2,3-dimethyl-5-phytylplastoquinol.

15 . . .

#### 6E. *Arabidopsis* Homologue to slr1737

- An *Arabidopsis* homologue to slr1737 was identified from a BLASTALL search using *Synechocystis* sp 6803 gene slr1737 as the query, in both public and proprietary databases. SEQ ID NO:109 and SEQ ID NO:110 are the DNA and translated amino acid sequences, respectively,  
20 of the *Arabidopsis* homologue to slr1737. The start is found at the ATG at base 56 in SEQ ID NO:109.

- The sequences obtained for the homologue from the proprietary database differs from the public database (F4D11.30, BAC AL022537), in having a start site 471 base pairs upstream of the start identified in the public sequence. A comparison of the public and proprietary sequences  
25 is provided in Figure 31. The correct start correlates within the public database sequence is at 12080, while the public sequence start is given as being at 11609.

Attempts to amplify a slr1737 homologue were unsuccessful using primers designed from the public database, while amplification of the gene was accomplished with primers obtained from SEQ ID NO:109.



Analysis of the protein sequence to identify transit peptide sequence predicted two potential cleavage sites, one between amino acids 48 and 49, and the other between amino acids 98 and 99.

#### 5 6F. slr1737 Protein Information

The slr1737 orf comprises 363 amino acid residues and has a predicted MW of 41kDa (SEQ ID NO: 39). Hydropathic analysis indicates the protein is hydrophillic (Figure 32).

10 The *Arabidopsis* homologue to slr1737 (SEQ ID xx) comprises 488 amino acid residues, has a predicted MW of 55kDa, and has a putative transit peptide sequence comprising the first 98 amino acids. The predicted MW of the mature form of the *Arabidopsis* homologue is 44kDa. The hydropathic plot for the *Arabidopsis* homologue also reveals that it is hydrophillic (Figure 33). Further blast analysis of the *Arabidopsis* homologue reveals limited sequence identity (25 % sequence identity) with the beta-subunit of respiratory nitrate reductase. Based on the sequence identity to nitrate reductase, it suggests the slr1737 orf is an enzyme that likely involves general  
15 acid catalysis mechanism.

Investigation of known enzymes involved in tocopherol metabolism indicated that the best candidate corresponding to the general acid mechanism is the tocopherol cyclase. There are many known examples of cyclases including, tocopherol cyclase, chalcone isomerase, lycopene cyclase, and aristolochene synthase. By further examination of the microscopic catalytic  
20 mechanism of phytoplastoquinol cyclization, as an example, chalcone isomerase has a catalytic mechanism most similar to tocopherol cyclase. (Figure 34).

Multiple sequence alignment was performed between slr1737, slr1737 *Arabidopsis* homologue and the *Arabidopsis* chalcone isomerase (Genbank:P41088) (Figure 35). 65% of the conserved residues among the three enzymes are strictly conserved within the known chalcone  
25 isomerases. The crystal structure of alfalfa chalcone isomerase has been solved (Jez, Joseph M., Bowman, Marianne E., Dixon, Richard A., and Noel, Joseph P. (2000) "Structure and mechanism of the evolutionarily unique plant enzyme chalcone isomerase". *Nature Structural Biology* 7: 786-791.) It has been demonstrated tyrosine (Y) 106 of the alfalfa chalcone isomerase serves as the general acid during cyclization reaction (Genbank: P28012). The

equivalent residue in slr1737 and the slr1737 *Arabidopsis* homolog is lysine (K), which is an excellent catalytic residue as general acid.

The information available from partial purification of tocopherol cyclase from *Chlorella protothecoides* (U.S. Patent No. 5,432,069), *i.e.*, described as being glycine rich, water soluble  
5 and with a predicted MW of 48-50kDa, is consistent with the protein informatics information obtained for the slr1737 and the *Arabidopsis* slr1737 homologue.

All publications and patent applications mentioned in this specification are indicative of the level of skill of those skilled in the art to which this invention pertains. All publications and  
10 patent applications are herein incorporated by reference to the same extent as if each individual publication or patent application was specifically and individually indicated to be incorporated by reference.

Although the foregoing invention has been described in some detail by way of illustration and example for purposes of clarity of understanding, it will be obvious that certain changes and  
15 modifications may be practiced within the scope of the appended claim.

## CLAIMS

What is claimed is:

1. An isolated nucleic acid sequence encoding a prenyltransferase.
- 5 2. An isolated nucleic acid sequence according to Claim 1, wherein said prenyltransferase is selected from the group consisting of straight chain prenyltransferase and aromatic prenyltransferase.
3. An isolated DNA sequence according to Claim 1, wherein said nucleic acid sequence is isolated from a eukaryotic cell source.
- 10 4. An isolated DNA sequence according to Claim 3, wherein said eukaryotic cell source is selected from the group consisting of mammalian, nematode, fungal, and plant cells.
5. The DNA encoding sequence of Claim 4 wherein said prenyltransferase protein is from *Arabidopsis*.
6. The DNA encoding sequence of Claim 5 wherein said prenyltransferase protein is encoded by a sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID  
15 NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:13, SEQ ID NO:14, SEQ ID NO:15, and SEQ ID NO:16.
7. The DNA encoding sequence of Claim 4 wherein said prenyltransferase protein is from soybean.
8. The DNA encoding sequence of Claim 7 wherein said prenyltransferase protein is encoded by  
20 a sequence comprising a nucleotide sequence selected from the group consisting of SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, and SEQ ID NO:23.
9. The DNA encoding sequence of Claim 7 wherein said prenyltransferase protein is encoded by a sequence selected from the group consisting of SEQ ID NO:95, and SEQ ID NO:96.
10. The DNA encoding sequence of Claim 7 wherein said prenyltransferase protein has an  
25 amino acid sequence selected from the group consisting of SEQ ID NO:97, and SEQ ID NO:98.
11. The DNA encoding sequence of Claim 4 wherein said prenyltransferase protein is from corn.
12. The DNA encoding sequence of Claim 11 wherein said prenyltransferase protein is encoded by a sequence comprising a nucleotide sequence selected from the group consisting of SEQ ID NO:25, SEQ ID NO:26, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:31, SEQ ID NO:104,  
30 SEQ ID NO:105, and SEQ ID NO:106.

13. The DNA encoding sequence of Claim 4 wherein said prenyltransferase protein is from rice.

14. The DNA encoding sequence of Claim 13 wherein said prenyltransferase protein is encoded by a sequence comprising SEQ ID NO:99.

5 15. The DNA encoding sequence of Claim 4 wherein said prenyltransferase protein is from wheat.

16. The DNA encoding sequence of Claim 15 wherein said prenyltransferase protein is encoded by a sequence comprising SEQ ID NO:100.

17. The DNA encoding sequence of Claim 4 wherein said prenyltransferase protein is from leek.

10 18. The DNA encoding sequence of Claim 17 wherein said prenyltransferase protein is encoded by a sequence comprising a nucleotide sequence selected from the group consisting of SEQ ID NO:101, and SEQ ID NO:102.

19. The DNA encoding sequence of Claim 4 wherein said prenyltransferase protein is from canola.

15 20. The DNA encoding sequence of Claim 19 wherein said prenyltransferase protein is encoded by a sequence comprising SEQ ID NO:103.

21. The DNA encoding sequence of Claim 4 wherein said prenyltransferase protein is from cotton.

22. The DNA encoding sequence of Claim 21 wherein said prenyltransferase protein is encoded by a sequence comprising SEQ ID NO:107.

20 23. The DNA encoding sequence of Claim 4 wherein said prenyltransferase protein is from tomato.

24. The DNA encoding sequence of Claim 23 wherein said prenyltransferase protein is encoded by a sequence comprising SEQ ID NO:108.

25 25. An isolated DNA sequence according to Claim 4, wherein said prokaryotic source is a *Synechocystis* sp.

26. A nucleic acid construct comprising as operably linked components, a transcriptional initiation region functional in a host cell, a nucleic acid sequence encoding a prenyltransferase, and a transcriptional termination region.

27. A nucleic acid construct according to Claim 26, wherein said nucleic acid sequence encoding prenyltransferase is obtained from an organism selected from the group consisting of a eukaryotic organism and a prokaryotic organism.

5 28. A nucleic acid construct according to Claim 27, wherein said nucleic acid sequence encoding prenyltransferase is obtained from a plant source.

29. A nucleic acid construct according to Claim 28, wherein said nucleic acid sequence encoding prenyltransferase is obtained from a source selected from the group consisting of *Arabidopsis*, soybean, corn, rice, wheat, leek canola, , leek, cotton, and tomato.

10 30. A nucleic acid construct according to Claim 26, wherein said nucleic acid sequence encoding prenyltransferase is obtained from a *Synechocystis* sp.

31. A plant cell comprising the construct of 26.

32. A plant comprising a cell of Claim 31.

33. A feed composition produced from a plant according to Claim 32.

34. A seed comprising a cell of Claim 31.

15 35. Oil obtained from a seed of Claim 34.

36. A natural tocopherol rich refined and deodorised oil which has been produced by a method of treating an oil according to Claim 35 by distilling under low pressure and high temperature, wherein said refined oil has reduced free fatty acids and a substantial percentage of tocopherol present in the pretreated oil.

20 37. A refined oil according to claim 36, wherein the pretreated oil is crude or pre-treated soybean oil.

38. A refined oil according to claim 36, wherein the refined oil is degummed and bleached.

25 40. A method for the alteration of the isoprenoid content in a host cell, said method comprising; transforming said host cell with a construct comprising as operably linked components, a transcriptional initiation region functional in a host cell, a nucleic acid sequence encoding prenyltransferase, and a transcriptional termination region,

wherein said isoprenoid compound selected from the group of tocopherols and tocotrienols .

30 41. The method according to Claim 40, wherein said host cell is selected from the group consisting of a prokaryotic cell and a eukaryotic cell.



42. The method according to Claim 41, wherein said prokaryotic cell is a *Synechocystis* sp.

43. The method according to Claim 41, wherein said eukaryotic cell is a plant cell.

44. The method according to Claim 43, wherein said plant cell is obtained from a plant selected from the group consisting of *Arabidopsis*, soybean, corn, rice, wheat, leek canola, , leek, cotton, and tomato.

45. A method for producing an isoprenoid compound of interest in a host cell, said method comprising obtaining a transformed host cell, said host cell having and expressing in its genome: a construct having a DNA sequence encoding a prenyltransferase operably linked to a transcriptional initiation region functional in a host cell,

wherein said prenyltransferase is involved in the synthesis of tocopherols,

and wherein said isoprenoid compound selected from the group of tocopherols and tocotrienols.

46. The method according to Claim 45, wherein said host cell is selected from the group consisting of a prokaryotic cell and a eukaryotic cell.

47. The method according to Claim 46, wherein said prokaryotic cell is a *Synechocystis* sp.

48. The method according to Claim 46, wherein said eukaryotic cell is a plant cell.

49. The method according to Claim 48, wherein said plant cell is obtained from a plant selected from the group consisting wherein said compound selected from the group of *Arabidopsis*, soybean, corn, rice, wheat, leek canola, , leek, cotton, and tomato.

50. A method for increasing the biosynthetic flux in a host cell toward production of an isoprenoid compound, said method comprising;

transforming said host cell with a construct comprising as operably linked components, a transcriptional initiation region functional in a host cell, a DNA encoding a prenyltransferase, and a transcriptional termination region,

wherein said isoprenoid compound selected from the group of tocopherols and tocotrienols,

51. The method according to Claim 50, wherein said host cell is selected from the group consisting of a prokaryotic cell and a eukaryotic cell.

52. The method according to Claim 51, wherein said prokaryotic cell is a *Synechocystis* sp.

53. The method according to Claim 51, wherein said eukaryotic cell is a plant cell.

54. The method according to Claim 50, wherein said plant cell is obtained from a plant selected from the group consisting *Arabidopsis*, soybean, corn, rice, wheat, leek canola, , leek, cotton, and tomato.

55. The method according to Claim 50, wherein said transcriptional initiation region is a seed-  
5 specific promoter.


[illegible]

**FIG. 1**

2/48


ATPT2	280	*	300	*	320	*	340	*	299
ATPT3									314
ATPT4									300
ATPT8									259
ATPT12									294
ATPT2									375
ATPT3									392
ATPT4									390
ATPT8									321
ATPT12									373
ATPT2	460	*	480	*					393
ATPT3									407
ATPT4									431
ATPT8									-
ATPT12									387

FIG. 1 (CONT)


**FIG. 2**




4/48

**FIG. 3**


5/48

**FIG. 4**


6/48

**FIG. 5**


7/48

**FIG. 6**


8/48

**FIG. 7**







10/48

**FIG. 9**


11/48

**FIG. 10**

12/48




13/48




**FIG. 12**




14/48

**FIG. 13**


15/48

**FIG. 14**


16/48

**FIG. 15**

17/48


**FIG. 16**

18/48


**FIG. 17**




19/48

**FIG. 18**

20/48

**FIG. 19**

21/48

**FIG. 20**

22/48

SLR1736 : PATIQAFWR-----ESRPHITIGTILS-----WAAVEITLGLD--G-NSWSPAS-----IDIFGAMTACIGNVIVGT : 64  
 SLR0926 : WVAQTPSS---PPLMLTIIYLRHKE-----AGRLIENPALWA--NCLAAQLPP-----IPUIGTIAIGTATSGLGCW : 68  
 SLL1899 : VTSUKIHRHDSMGVCKSYQLTKE-----RIPILITTTAAS--WTASEPRVD-----IPKILITIGGTAAASAQT : 71  
 SLR0056 : PSDIQNTGO--NQAKQROLLOKGAAGCESSINKURLQKPETWIPLWGVCEAASSGGVINSVEDFKAUTCMHSGPMTGTQT : 88  
 SLR1518 : TESSPLAP--STAPTRKLMAAIAK-----PAYTAATVGSNAAYGIGTQWH-----GQFTIFISAFALIAINLS : 71

M P 6 9 1 L

SLR1736 : NOE3VDITRINKPNLPANGDEFAUGRNIGCCGASALWIGIGLGLTVGIS--LIGTAYSPPRRLKRFSLLAALCETVRG : 152  
 SLR0926 : NQETRDLUPQVETKORPLAARAI-SVQVCGALMA--HQAQIAFHTPLS-----FWCVAAVPRVAYPGA RVFPVQVL-S : 150  
 SLL1899 : NQDIDQVYELRTRARIPAGKQ-QPRHAIIFADALEHSEFALIAIENNVLSGC---LALSIVFYLYTHWLKRHTAQNIG- : 156  
 SLR0056 : NQFYLDQDAINEPYRPSGHSISPPQVTCILELWAGGVYQIDVQAQHPPIMMVTIGSAFNAVYSAPPLKLNKONGWGNV- : 177  
 SLR1518 : NQVETLSGTGIDVRKHSVNLGTGRNLVELISNEFTIASLGLASISRAQDK-----VDEISVAFHGYTYQGPFRUGVLCGELIC : 157

N 50 Did 1 a 6 6 1 6

SLR1736 : VVNLGLELIRIGLGYPPITTPVWVITFELHFTVAIAIKDVPMQGRQFKIQELIOTIKONVFRGTALTLG-CYDAMAWG : 241  
 SLR0926 : AWGAVNHSAS---HNGDITDATWVINGATVFMIGEDTVYVADREEDRHSNSALFFI-QYAGEAGCFEA--ITGCLFY : 234  
 SLL1899 : AGSIPPLGCA-----AVTGDLSWTPWVIFALHFMPPHFWALLAMIKDDVAGVNPMPPTALAEKTVSQWYYS-- : 241  
 SLR0056 : GASVIAIPWAG-HAIFGTINPTIMVHLYSLAGLGAVVNDFKSAPQROHKKSPHMF-IGTAAWICAMT--DVFQAG : 263  
 SLR1518 : LITGCPUIAIAVYYSQSESHNLTPTSPVFGSIAHLCFCHFHVQVDELLAAGKKPIRLI-TQSGSALTSIVSV : 246

1 1 vl t e d 6 6

FIG. 21

23/48

	280	*	300	*	320	*	340
SKR1736 :	AMPLNTA <sup>N</sup> IN <sup>N</sup> SH <sup>N</sup> CH <sup>N</sup> LAI <sup>N</sup>	WNRSRDVHLESKTE <sup>N</sup>	ASFVQ <sup>N</sup> IN <sup>N</sup> KI <sup>N</sup>	FLEV <sup>N</sup> LY <sup>N</sup> PT <sup>N</sup> AP <sup>N</sup> WPNESNTIF <sup>N</sup>			: 308
SLR0926 :	LM <sup>N</sup> NP <sup>N</sup> Y <sup>N</sup> NI <sup>N</sup> SLA <sup>N</sup>	AGW <sup>N</sup> IOY <sup>N</sup> IQ <sup>N</sup> LSAP <sup>N</sup> TEPKLYQ <sup>N</sup> --	IGQ <sup>N</sup> NV <sup>N</sup> I <sup>N</sup> IG <sup>N</sup> CV <sup>N</sup> LA <sup>N</sup>	LI <sup>N</sup> GW <sup>N</sup>			: 292
SLL1899 :	LHOLG <sup>N</sup> LAI <sup>N</sup> AI <sup>N</sup> IGG <sup>N</sup>	EVKAWOLKQ <sup>N</sup> GD <sup>N</sup> RG <sup>N</sup> ARG <sup>N</sup> --	LEKES <sup>N</sup> LY <sup>N</sup> IM <sup>N</sup> ICLA <sup>N</sup>	VIDSPVTHOLVQMG <sup>N</sup> TLIG <sup>N</sup>			: 316
SLR0056 :	Y <sup>N</sup> HQO <sup>N</sup> YI <sup>N</sup> AT <sup>N</sup> IL <sup>N</sup>	PE <sup>N</sup> TFQ <sup>N</sup> DMY <sup>N</sup> FLRM <sup>N</sup> LEND <sup>N</sup> YKYO <sup>N</sup> --	ASAQ <sup>N</sup> PE <sup>N</sup> LV <sup>N</sup> FC <sup>N</sup> AT <sup>N</sup> GL <sup>N</sup> GHAGI <sup>N</sup>				: 324
SLR1518 :	QAPWOTLL <sup>N</sup> EP <sup>N</sup> AS <sup>N</sup> IP <sup>N</sup> WAV <sup>N</sup>	FRHVGOYHDO <sup>N</sup> EQVSNCKE <sup>N</sup> IAVNLH <sup>N</sup> ES <sup>N</sup>	AGY <sup>N</sup> GHAG <sup>N</sup> IG <sup>N</sup>				: 307

**FIG. 21 (CONT.)**



24/48

```

ATPT2 : -----MESLSSSSLSAAGFCWKKQNLKLSLSEIRVLRCDSSKVAKPKFRNNLVRPDGCGSSLLLPKHKSFRVFNATAG : 80
SLR1736 : -----
ATPT3 : MAFFGLSRVSRRLKSSVSPTSSSALLQSOHKSLSNPVTTHYTNPETKCPWSNDNYQVHNSKGRHLEHKEFGVGNRYRLICGMSS : 89
SLR0926 : -----
ATPT4 : -----MWRRSWYRESSRISVSSSLPNRLIPWSRELCAVNSQ?-----PVSTESTAKLGITGVRSDANRVEATA : 67
SLR1899 : -----MVT : 4
ATPT12 : -----MTSILATVSTIHSRVTSVDRVGVLSLRNSDSVEFT-----RRSGFSTLIESPGRREVVRRAETDT : 63
SLR0056 : -----MSDT : 4
ATPT8 : -----MVL : 4
SLR1518 : -----MTES : 4

```

```

ATPT2 : PEAFDSNSKOK-----SFRSDDAFYR-----FSRPHIGTIVLSLS-----VSELAENKVS--DISPLFTGLE : 140
SLR1736 : -----MATHQAEWR-----FSRPHIGTIVLSLS-----LLTILGDN-SVNSPASJOLYEG : 49
ATPT3 : SSVLEGPCKKODKEKSGVWVKASWDLYLPEVRGYAKLARLDKPIGTWILAWPCMS-----IALADPS--LPSFKYALFGC : 170
SLR0926 : -----MVAQTPSP-----PLMTIIL-----LRHHPAGRIIMIPALMA-----VCLAO--G--LPPLDICTAL : 56
ATPT4 : TAAATATATTG-----EISSRVAALAGIGHHYAR-----CYWELSKANISMLVWATSG-----TGYILGTQNAALISFPGCYTCAG : 138
SLR1899 : TKIHRHDSMG-----AVCKSYQLTKP-----RIIPLIITTAASMI-----ASER--VOLPKHITELG : 60
ATPT12 : DKVKSQTPDKAP-----AGGSINOLLCKGAS-----QETNKWKIRIOLTKPVTIPLVWGWCCGAASGNEHNTPEIAKSLC : 139
SLR0056 : QNT-GQNOAKA-----ROLLCKGAAP-----GESSINWKIRIQIWKPIPIPLINGWCCGAASSGYINSVEDEIKATC : 73
ATPT8 : EVPKLASAAEY-----FFKRGQCKQF-----RSTILIPATAINLRVP-----EALIGEST--DIVTSEHRVOR : 63
SLR1518 : SPLAPSTAPAT-----RKLMAAIAKP-----PHYTAVVVPIIVG-----SAGYGLTG--QNHGQFTIFLL : 59

```

FIG. 22

25/48

80 : AVAAALANIVYVIGMQLSVETDKKVKPYLPLASGEYSNTGLAVASESEFWLGMVGSHPLEWALPSEFGLGTAYS-INIPALR : 228  
 SLR1736 : AMIACLLGNVYVIGMQLSVETDKKVKPYLPLASGEYSNTGLAVASESEFWLGMVGSHPLEWALPSEFGLGTAYS-INIPALR : 134  
 ATPT3 : GALL---KRGAGCTIDILEQDITKVDITKLRKIASGLIT-PTQCGELGLOUELGIG-----ILLQNNYSYVAGAS-----SHLAVF : 246  
 SLR0926 : GTIA---TSGLCGVADIMVADIRKIDPOVERTKQRLAARAS-VQVCGVAVFALICAG-----IAFYLTPISTFVACV-----AVPQIV : 132  
 ATPT4 : TMGI-----AASANSACVHEISNE SKMKRTMLRLPSGRISVPHAVAWATAGAGACL-----LESKTNMAAGASAN-----IULYAF : 215  
 SLR1899 : GTIA---AASACTIDILEQDITKVDITKLRKIASGLIT-PTQCGELGLOUELGIG-----ILLQNNYSYVAGAS-----SHLAVF : 137  
 ATPT12 : MNSGPGTGYTOTINDMVRDIAINEPYRPIESGAISEPEVITQVWVILGCGIGLIGILD-VWAGHTTPTVYALG-----GSTLSY : 223  
 SLR0056 : MNSGPGTGYTOTINDMVRDIAINEPYRPIESGAISEPEVITQVWVILGCGIGLIGILD-VWAGHTTPTVYALG-----GSTLSY : 157  
 ATPT8 : GIAE---ITEMIHVASLTHEDVINDADTRGVGSLNVVGNKMSVAGDFISRACGL-----ALKNTERVAUJATAVEHLVTGETM : 144  
 SLR1518 : SADA---TIAHINLSDDITESTGIDVRAHSVVNLTGVRNLVELSNFFELACULGMSMS--WRAQDWTALELIGV-----IFEGY : 138

n d d

FIG. 22 (CONT. -1)

26/48

ATPT2 : WKR-FALVAMCILAARAIIVQIAEYTH-IQTHVFGRIPIETRPTEPATAEMSPFESVIALEKIDIPDIQKI-ICRPSFTIC-QKR : 313  
 SLR1736 : LKR-ESLLAALCILTRGIVNGLGIF-ERIGLGYPTIITPTM-ETILFIMETIAIAFKDVPDMEGRQ-FKQCTITIQIE-KQN : 218  
 ATPT3 : SYP-LMKRETFWQAFGLTINWGAJG--IT--AKGSIAPSKTPIYLSGVCIVYDTIYAHODKEDVK-VKSKSTAIRFE-DNT : 328  
 SLR0926 : AYP-GAKRVFVPQILSLIANGFAMIS--S--ATSDIDATM--WQATVETUIGFDTVYAHADRDERR-HQVNSAIFFE-QYV : 213  
 ATPT4 : VET-PLKQLHPINTWCAVGAIPPIG--A--ASSQISYNSI--IPAALYFQIPHEALAHLCRNDYAA-GEYKMSIFEDP---S : 294  
 SLR1899 : VETHMLKRHTAQNIIVCGAAGSIPPIG--A--ATQDISWTPM--IFALIFLKEPHEWALAMIKDDYAO-VNYPMPPIAIEEKT : 220  
 ATPT12 : IYS-APPLKQKQNGNFAICASYSLPMMAGQAFETLEPDVY--ITLLYSIAGEIAEVNDEKSVGGRA-DEKSLPAFFE-TET : 308  
 SLR0056 : IIS-APPLKQKQNGNFAICASYSLPMMAGQAFETLEPDVY--ITLLYSIAGEIAEVNDEKSVGGRA-DEKSLPAFFE-IGT : 242  
 ATPT8 : EITSSTEQRYSMDYVOKTYKKTASHINSCKAVAJLTCQAEVAMLAFFEYGRNLCEAFQIDDDIDFTGTSAIKCSIESDIRH--GV : 231  
 SLR1518 : TYGPPFERLGLGELICLITFGPRAL-AAAYSSQSFSFNNET-PSVTVGISATILFCSHFGQVDELA-AGKSPIRUE-TKL : 223

9 9

ATPT2 : VFTCVTLIQMAYAAIVGATSPFINSKVISVGHVIEATTEWARAKSVDLSSKTETS---CYNQIWKIFYAEYIHLPELK----- : 393  
 SLR1736 : VERCTELITGCYLMAHNGIWAAMPLNTAFITVSHLCILAEHWRSDVHLESKTEBAS---FYQIWKIFELEYEYIYIAJWLPNES : 304  
 ATPT3 : KLMITGFGTASIGFALSIGESADLGMQYASIAASGQGWQGTADLSSGADCS-----RKFSYKNWEGAI-FSCVVGCRSFO- : 407  
 SLR0926 : GEAVGIEFFALTIGCJFYTGMLMLNPLWLSFAIAI--VGMVIOYIQLSAPTEP-KHY-----GOIFGQNVIIIGFVIEBACILGWL--- : 292  
 ATPT4 : GKRPAAVAIRNCFYNIPIGELAYDHWGLTSSWECESTLITLAAATAESFYRDMHKA-----RKMHHASLFLPVMSCILHRVSN : 379  
 SLR1899 : VSQHWYYSILVVPESLIVYPLHQLGILYLAIAI--GGQFIVKAWQLQAPGDRDIA-----RGLNFSLEYLMLJCLASVIDSLPVT : 303  
 ATPT12 : AKVHCYGAIDITOLSVAGYLJASCKPYVALAJVALI--APQIFQFKVELKDPVKYDYK-----YOASAQPELVLGIEYTAIASOH---- : 387  
 SLR0056 : FAWFCYIMDOVQAGIAGVLYVHQQLVATIMTLI--IPQITFODMVELRNPLENDEYK-----YOASAQPELVGEGMELATGIAIGHAGI- : 324  
 ATPT8 : ITAPIIEAEFEFPQIREHVDVEKOPRNVDIAEYIGSDGFORARELAWEHANLAAAIIGSLPETDNEQKRSRRRAIIDDITHRVITRN : 320  
 SLR1518 : GSOVLTLSVSLYLITANGVICHQAPWOTLL-BASIPWAVOIRHVQVHDQPEQVSNCK----FIAVNLHFFSCMLMAAGYWGAGIG-- : 307

FIG. 22 (CONT. -2)

27/48

	*	460	*	480	*
ATPT2	:	-----	:	-----	:
SLR1736	:	NTIF-----	:	308	:
ATPT3	:	-----	:	-----	:
SLR0926	:	-----	:	-----	:
ATPT4	:	NOOQVVEAGLTNSVSGEVKTORRKKRVAQPPVAYASAPPELPAPSEVSP	:	431	:
SL11899	:	---HQLVAQMGTLILG---	:	316	:
ATPT12	:	-----	:	-----	:
SLR0056	:	-----	:	-----	:
ATPT8	:	K-----	:	-----	:
SLR1518	:	-----	:	-----	:

**FIG. 22 (CONT. -3)**





FIG. 23

29/48

**FIG. 24**



30/48




FIG. 25

31/48






FIG. 26

32/48






**FIG. 27**



33/48



**FIG. 28**



**FIG. 29A****FIG. 29B****FIG. 29C**

35/48

**FIG. 30A****FIG. 30B**



36/48

**FIG. 30C****FIG. 30D**

37/48

Query Sequence: F4D11 AL022537

Database: PIR\_T04448.atcea.list.fasta

Database: PIR\_T04448

Plus (+) denotes forward strand, and minus (-) reverse strand.

Asterisks (\*) denote bases not shown on pair wise alignments.

Alignment 1

```

Query-      12194 CACACGTTCTCGTCCTTTTCTTCTTCTCTGCAATTCTTCACAGAGTTTGTCAACCACCA
genomic
ATCEA4C371+  1 ----- C est
MET
Query-      12134 ACACCAACAGACAAATTTCACATTGTTTTCATTTTCGTTTCTTCTTCGATTATGGA
ATCEA4C371+  2 ACCCAAACATCACAATTTACATTCTTTGCATATTCTTCTTCTTCCATTATGGA

Query-      12075 GATACGGAGCTTGATTGTTTCTATGAACCTAATTATCTTCCTTTGAGCTCTCTCGCCC
ATCEA4C371+  62 GATACGGAGCTTGATTGTTTCTATGAACCTAATTATCTTCCTTTGAGCTCTCTCGCCC

Query-      12015 TGTATCTCCTCTCACTCGCTCACTAGTTCGGTTCGGATCGACTAACTAGTTCCTCGCTC
ATCEA4C371+  122 TGTATCTCCTCTCACTCGCTCACTAGTTCGGTTCGGATCGACTAACTAGTTCCTCGCTC

Query-      11955 CATTCTAGGGTTTCGGCGTTCATCTCCACCCGAATAGTGAACTGACAAGATCTCCGT
ATCEA4C371+  182 CATTCTAGGGTTTCGGCGTTCGATCTCCACCCGAATAGTGAACTGACAAGATCTCCGT

Query-      11895 TAAACCTGTTTACGTCCCGACGTCTCCCAATCGCGAACTCCGGACTGCTCAGAGTGGTA
ATCEA4C371+  242 TAAACCTGTTTACGTCCCGACGTCTCCCAATCGCGAACTCCGGACTGCTCAGAGTGG
Synecho seq aligns from
here

Query-      11835 AATTGATCCATTCCATTCCATTCTCTTCTTGTGTTTATTAAAGCTCCAATTCAG
ATCEA4C371+  299

```

FIG. 31

38/48

-- 60 bp removed --

Query- 11715 .....TTTG

ATCEA4C371+ 299

PIR:T04448 1

Query- 11655 GTGGCTCACCATTGACGACTACTTTTGAATTTGAGTTTTGAAAAATGCAATTTAACAT

ATCEA4C371+ 299

PIR:T04448 1

M Q F N I

arab sequence which is incorrect

Query- 11595 CAGAGAGTTTTTTTATGGTTGATAACTTATTGTTTAACTTTGAAAAATGCAG

ATCEA4C371+ 299

PIR:T04448 6

R E F F F L W L I T Y C L T F E K C R Y

Query- 11535 CCATTTTCGATGGAACACCTCGGAAGTTCTTCGAGGGATGGTATTTTCAGGGTTCCATCCC

ATCEA4C371+ 302 CCATTTTCGATGGAACACCTCGGAAGTTCTTCGAGGGATGGTATTTTCAGGGTTCCATCCC

PIR:T04448 26

H F D G T P R K F F E G W Y F S I P

Query- 11475 AGAGAAGAGGGAGAGTTTTGTTTTATGTATTCTGTGGAGAATCCTGCATTTCCGCAGAG

ATCEA4C371+ 362 AGAGAAGAGGGAGAGTTTTGTTTTATGTATTCTGTGGAGAATCCTGCATTTCCGCAGAG

PIR:T04448 46

E K R E S F C F M Y S V E N P A F R Q S

Query- 11415 TTTGTCACCATTGGAAGTGGCTCTATATGGACCTAGATTCAC TGGTGTGGAGCTCAGAT

ATCEA4C371+ 422 TTTGTCACCATTGGAAGTGGCTCTATATGGACCTAGATTCAC TGGTGTGGAGCTCAGAT

PIR:T04448 66

L S P L E V A L Y G P R F T G V G A Q I

**FIG. 31 (CONT-1)**

SUBSTITUTE SHEET (RULE 26)

39/48

```

Query- 11355 TCTTGGCGCTAATGATAAAATTTATGCCAATACGAACAAGACTCTACAATTTCTGGGG
          |||||||||||||||||||||||||||||||||||||||||||||||||||||-----
ATCEA4C371+ 482 TCTTGGCGCTAATGATAAAATTTATGCCAATACGAACAAGACTCTACAATTTCT
          ::::::::::::::::::::::::::::::::::::::::::::::::::::::::::::
PIR:T04448 86 L G A N D K Y L C Q Y E Q D S H N F W G
ATCEA4C371+ Exon 11538 11301 Confidence: 100 100

```

```

Query-      11295 AGGTAACCTCCTTGACCCTTAAATGCTGTGTCATGACAATAAGAAATCATATCTGAGTCT
              -----
ATCEA4C371+ 537
              :
PIR:T04448   106 D
PIR:T04448   Exon      11609      11294      Confidence: 100 100

```

```
Query-      11235 TTTCTCTACTTCTAGTACTAATGTTTCGTTATTGTTGTTAAAGATCTAAGTCTTATCTGAA
              -----
PIR:T04448   107
```

Query- 11175 TTTTGTTACATTTTGGTTCTGGTGCTTCTCAACATGAATTTGTATATATGACTTTAAAG  
-----  
PIR:T04448 107

```

Query-      11115  ATTGCTTACCTAAAGTTTTTACTCATGCATAGATCGACATGAGCTAGTTTTGGGGAATAC
                -----
PIR:T04448   107                                     R H E L V L G N T

```

```

Query-      11055 TTTTAGTGTCTGTGCCAGGCGCAAAGGCTCCAAACAAGGAGGTTCCACCAGAGGTTCTCAC
              ::::::::::::::::::::::::::::::::::::::::::::::::::::::::::::
PIR:T04448   116  F  S  A  V  P  G  A  K  A  P  N  K  E  V  P  P  E
PIR:T04448      Exon      11083      11004 Confidence: 96 100

```

```
Query-      10995 TCCTCCCTTGTGTTACTTTGTTATCTGTTAAATAGTTTTCCAATTGTATCCGGATAGT
              -----
PIR:T04448   133
```

Query- 10935 GTTCTACTTCTCCTTGTAGAAATCTCAAGTTTTGTTACTCTTGCTATTCTCTTGGATG  
-----  
PIR:T04448 133

**FIG. 31 (CONT-2)**

40/48

Query- 10875 TTGATTTGTAAAGCATGTCGTTTTATTGTAGGAATTTAACAGAAGAGTGTCCGAAGGGTT  
-----  
PIR:T04448 133 E F N R R V S E G F

Query- 10815 CCAAGCTACTCCATTTTGGCATCAAGGTCACATTTGCGATGATGGCCGGTAATTATATGA  
-----  
PIR:T04448 143 Q A T P F W H Q G H I C D D G R  
PIR:T04448 Exon 10844 10768 Confidence: 100 100

Query- 10755 TTCTATGCACAACAAGAATTCACATATTATAAATATTGGATATTGAGTATTTTGTGA  
-----  
PIR:T04448 159

Query- 10695 AAATTTCTGTGTTTAAATCTGACTTGACTTGTTTTGTCAGTACTGACTATGCGGAACTG  
-----  
PIR:T04448 159 T D Y A E T V

Query- 10635 TGAAATCTGCTCGTTGGGAGTATAGTACTCGTCCCGTTTACGGTTGGGGTGATGTTGGGG  
-----  
PIR:T04448 166 K S A R W E Y S T R P V Y G W G D V G A

Query- 10575 CCAAACAGAAGTCAACTGCAGGCTGGCCTGCAGCTTTTCCTGTATTGAGCCTCATTGGC  
-----  
PIR:T04448 186 K Q K S T A G W P A A F P V F E P H W Q

Query- 10515 AGATATGCATGGCAGGAGGCCTTTCCACAGGTGTGAGCTTGCTTGATTGACTTAAAGTT  
-----  
PIR:T04448 206 I C M A G G L S T G  
PIR:T04448 Exon 10655 10486 Confidence: 96 100

Query- 10455 AATAAATAGACGGTTAAGTTTACTTGCTAGTACTAACAGAAAATTAAGAAAGAAACCAC  
-----  
PIR:T04448 216

**FIG. 31 (CONT-3)**

SUBSTITUTE SHEET (RULE 26)

41/48

Query- 10395 CCTCTTCTATCAGCAGAACTGCTATTGTAGTTCTTATTTTCTCTTGTATTGCAGG  
-----  
PIR:T04448 216

Query- 10335 GTGGATAGAATGGGGCGGTGAAAGGTTGAGTTTCGGGATGCACCTTCTTATTCAGAGAA  
.....  
PIR:T04448 216 W I E W G G E R F E F R D A P S Y S E K

Query- 10275 GAATTGGGGTGGAGGCTTCCCAAGAAAATGGTTTGGGTAAAACATTCATCCTTTTGCT  
.....  
PIR:T04448 236 N W G G G F P R K W F W  
PIR:T04448 Exon 10336 10239 Confidence: 96 100

Query- 10215 ACATTCTTGTTCAGACTTTAGTTAGCTAGTGGACCTGTGTATACACCCACATATAGTA  
-----  
PIR:T04448 248

Query- 10155 TACTTGTTTGATAGCTTTATTTGTCAATGTCTCTTTACAGGTCCAGTGAATGTCTTTGA  
.....  
PIR:T04448 248 V Q C N V F E

Query- 10095 AGGGGCAACTGGAGAAGTTGCTTTAACCGCAGGTGGCGGGTTGAGGCAATTGCCTGGATT  
.....  
PIR:T04448 255 G A T G E V A L T A G G G L R Q L P G L

Query- 10035 GACTGAGACCTATGAAAATGCTGCACTGGTATGCACTTATAAGATCTTCTTAAGCAATGA  
.....  
PIR:T04448 275 T E T Y E N A A L  
PIR:T04448 Exon 10115 10008 Confidence: 100 100

Query- 9975 CAGTGAGTATTAGAAGGCAGATAGTTTACAAAAGCTCTGGGCCCTTGTAATCTGCAGGT  
-----  
PIR:T04448 284 V

Query- 9915 TTGTGTACACTATGATGGAAAATGTACGAGTTTGTTCCTTGAATGGTCTTGTAGATG  
.....  
PIR:T04448 285 C V H Y D G K M Y E F V P W N G V V R W  
-----  
GSDS:S:495- 532 tagatg

**FIG. 31 (CONT-4)**

SUBSTITUTE SHEET (RULE 26)



42/48

```
Query-      9855 GGAAATGTCTCCCTGGGG TTATTGGTATATAACTGCAGAGAACGAAAACCATGTGGTAA
          :-----:
PIR:T04448   305 E M S P W G Y W Y I T A E N E N H V
          |||||---|||||---|||||
GSDB:S:495-  526 ggaaat tctccctgggggttattggtatataactgcagagaNcgNaaaccatgtg
PIR:T04448    Exon       9917       9801 Confidence: 100 100
GSDB:S:495-    Exon       9961       9801 Confidence:  93  93

Query-      9796 ATTTGTTTTACTAGTTTCATTCACTTTTACTTTTGACATCATATCATTCCCCTTATGGCTA
          -----
PIR:T04448   323
          -----
GSDB:S:495-  471

Query-      9736 GATTCCAACACCCGATGAATGTCTTGTGACAGGTGGAAGTAGAGGCAAGAACAAATGAAG
          -----:-----
PIR:T04448   323                               V E L E A R T N E A
          -----:-----
GSDB:S:495-  471                               gtggaactagaggcNagaacaaatgaag

Query-      9676 CGGGTACACCTCTGCGTGCTCTACCACAGAAGTTGGGCTAGCTACGGCTTGCAGAGATA
          :-----:
PIR:T04448   333 G T P L R A P T T E V G L A T A C R D S
          |||||
GSDB:S:495-  443 cgggtacacctctgcgtgctcctaccacagaagltgggctagctacggcttgcaagata

Query-      9616 GTTGTACGGTGAATTGAAGTGCAGATATGGGAACGGCTATATGATGGAAGTAAAGGCA
          :-----:
PIR:T04448   353 C Y G E L K L Q I W E R L Y D G S K G K
          |||||
GSDB:S:495-  383 gttgttacggtgaattgaagtgcagatatgggaacggctatatgatggaagtaaaggca

Query-      9556 AGGTATGTATGCTAATGTGATCCAATCCCTGTAGTTAAAAGTCTTAACAAATCCTAAGGC
          ::-----:
PIR:T04448   373                               L K V L T N P K A
          ||-----
GSDB:S:495-  323 ag
PIR:T04448    Exon       9704       9555 Confidence: 100 100
GSDB:S:495-    Exon       9704       9555 Confidence:  98 100
```

**FIG. 31 (CONT-5)**

```

Query-          9496 AGTGAAGAAGATTATGAACGTTTGTATGGTTAACAATGATGCAGGTGATATTAGAGAC
                  ::::::::::::::::::::::::::::::::::::::::::::::::::::::::::::::::::::
PIR:T04448      382  V K E D Y E R L L W L T M M Q V I L E T
                  -----|||
GSDB:S:495-     321                                     gtgatattagagac

```

```

Query-          9376 AGATACGAGCAACACGCCCCGAGCTACTAAACAGGCTCTTCAGGTCCCATTGGATCTTGA
                  ::::::::::::::::::::::::::::::::::::::::::::::::::::::::::::::
PIR:T04448      422  D T S N T P E L L K Q A L Q V P L D L E
                  |||
GSDB:S:495-     247 agatacgaacacgccccgagctactaaaacaggctcttcagggtcccattggatcttga

```

Query- 9316 AAGCGCCTTAGGTTTGGTCCCTTTCTTCAAGCCACCGGTCTGTAAGATTGATGAGTGCT  
(stop)  
PIR:T04448 442 S A L G L V P F F K P P G L  
|||||  
GSDB:S:495- 187 aagcgcccttaggtttgggtccctttcttcaagccaccgggtctgtaacattgatgagtgtt  
PIR:T04448 Exon 9522 9274 Confidence: 100-100

Query- 9256 ~~TTCATTTGAGACCACTCAGCATTAACCTTATTCTATCGTAG~~  
-----  
PIR:T04448 456  
|||||  
GSDB:S:495- 127 ttgtttggtgatagaccatgtgatgaatgaagccttagtcattgccttagcttc

```

Query-          9196 ACTATTATGTATGTATGATTTTAGTTTCGTTCCGTCCTTGTGGTAAATGATACGGGCCAGT
                  |||
GSDB:$:495-     67 actattatgtatgtatgatlttagtttcgttccggtccttgtggtaaatgatacggggcagt

```

```

Query-      9136 GTAAAGTCTAGTTCAATAAAAGCCTTGAGTCGCATAATTTCAATTTCAAATTGCATC
              |||||-----
GSDB:S:495-  7 gtaaagt
GSDB:S:495-  Exon      9450      9130  Confidence: 98 100

```

**FIG. 31 (CONT-6)**

44/48


ATCEA4C37145\_1 3063693/emb|CAA18584.1| 4.0e-43 (AL022537) putative protein  
[Arabidopsis thaliana]

PIR:T04448 sPIR-T04448 shypothetical protein F4D11.30 - Arabidopsis thaliana;  
g3063693|emb|CAA18584.1 (AL022537) putative protein [Arabidopsis thaliana]\_F4D11.30


GSDB:S:4955486|AI995392|AI995392|701673779 A. thaliana, Columbia Col-0,  
inflorescence-1 Arabidopsis thaliana cDNA clone 701673779, mRNA sequence.


**FIG. 31 (CONT-7)**

45/48

**FIG. 32**

46/48

**FIG. 33**



**FIG. 34**



48/48

slr1737 SYNSP S74814  
slr1737 ARATH T04448  
CFI ARATH P41088

-----M  
MEIRSLIVSMNPNLSSFELSRPVSPLTRSLVPFRSTKLVPRSISRVSASI  
-----

slr1737 SYNSP S74814  
slr1737 ARATH T04448  
CFI ARATH P41088

KFP-----PHSGYHWQGS-PFEGWYVRL  
STPNSETDKISVKPVYVPTSPNRELRTPHSGYHFDGTPRKPFEGWYFRVS  
-----

slr1737 SYNSP S74814  
slr1737 ARATH T04448  
CFI ARATH P41088

LPQSGESFAFMYSIENPASDHHYGGGAVQILGPATK----KQENQEDQLV  
IPEKRESFCFMYSVENPAFRQSLSPLEVALYGPRFTGVGAQILGANDKYL  
MSSSNACASPSPPFA----VTKLHVDVS-

slr1737 SYNSP S74814  
slr1737 ARATH T04448  
CFI ARATH P41088

WRTFPSVKKFWASPRQFALG-HWGKCRDNQ-AKPLLSEEFFATVKEGYQ  
CQYEQDSHNFWDGRHVLGNTFSAPVGAAPNKEVPPEEFNRRVSEGFQ  
--TFVPSVKSPASSNPLELG-GAGVRGLDIQ-GK-----FVIFTVIGVY

slr1737 SYNSP S74814  
slr1737 ARATH T04448  
CFI ARATH P41088

IHQNHQGGQIIHGDR-----HCRWQFTVEPEVTWGSPNRFPRATAGW  
ATPFWHQGHICDDGRDYAETVKSARWEYSTRPVYWGVDGAKQKSTAGW  
LEGNAPVPSLSV-----KWKGKTTEELTESIPFFREIVTGAF

slr1737 SYNSP S74814  
slr1737 ARATH T04448  
CFI ARATH P41088

LSFLPLFDPGWQILLAQGRAHWLKWQREQYEFDHVLYAEKNWGHSPS  
PAAFPVFEPHWQICMAGGLSTGWIEWGERFEFRDAPSYSEKNWGGGFPR  
EKFIKVT-----M-----KLPLTGQYSEKVTENC

slr1737 SYNSP S74814  
slr1737 ARATH T04448  
CFI ARATH P41088

RWFVLQANYFPDHPG-LSVTAAGGERIVLGRPE---EVALIGLHHQGNFY  
KWEVWQCNVFEAGTGEVALTAGGGLRQLPGLTETYENAALVCVHYDGMY  
VAIWKQLGLYTDCEA-KAV----EKFLIFKE---ET-----

slr1737 SYNSP S74814  
slr1737 ARATH T04448  
CFI ARATH P41088

EFPGHGTVTWQVAPWGRWQLKASNDRYWVKLSGKTDKKGSLVHTP-TAQ  
EFVPWNGVVRWEMSPWGYWYITAENENHVVELEARTNEAGTPLRAPTEV  
-FPPG-SSILFALSPTGSLTVAFSKDDS-IPETGIAVIENKLLAEA-VLE

slr1737 SYNSP S74814  
slr1737 ARATH T04448  
CFI ARATH P41088

GLQLNCRDTRGYLYLQLGSGVGHG-----LIVQGETDTAGLEVGG-----  
GLATACRDSYGELKLQIWERLYDGSKGSVILETKSSMAAVEIGGGPWFG  
--SIIGKNGVSPGTRLSVAERLSQ-----LMMKNKDEKEVSDHSL-----

slr1737 SYNSP S74814  
slr1737 ARATH T04448  
CFI ARATH P41088

----DWGLTEENLSKKT-----VPF-----  
TWKGDTSNTPELLKQALQVPLDLESALGLVPFFKPPGL  
----EEKLAKEN-----

FIG. 35

## SEQUENCE LISTING

<110> Lassner, Michael  
Post-Beittenmiller, Martha  
Savidge, Beth  
Weiss, James

<120> Nucleic Acid Sequences Involved in  
Tocopherol Synthesis

<130> 17133/00/WO

<150> 60/129,899

<151> 1999-04-15

<150> 60/146,461

<151> 1999-07-30

<150> PCT/US00/10368

<151> 2000-04-14

<160> 94

<170> FastSEQ for Windows Version 4.0

<210> 1

<211> 1182

<212> DNA

<213> Arabidopsis sp

<400> 1

atggagtctc tgctctctag ttcttctctt gtttccgctg ctggtgggtt ttgttggaag

60

aagcagaatc taaagctcca ctctttatca gaaatccgag ttctgcgttg tgattcgagt

120

aaagttgtcg caaaaccgaa gtttaggaac aatcttggtta ggcctgatgg tcaaggatct

180

tcattgttgt tgtatccaaa acataagtcg agatttcggg ttaatgccac tgcgggtcag

240

cctgaggctt tcgactcgaa tagcaaacag aagtctttta gagactcgtt agatgcgttt

300

tacaggtttt ctaggcctca tacagttatt ggcacagtgc ttagcatttt atctgtatct

360

ttcttagcag tagagaaggt ttctgatata tctcctttac ttttcaactgg catcttgag

420

gctgttgttg cagctctcat gatgaacatt tacatagttg ggctaaatca gttgtctgat

480

gttgaaatag ataagggttaa caagccctat ctccattgg catcaggaga atattctgtt

540

aacaccggca ttgcaatagt agcttccttc tccatcatga gtttctggct tgggtggatt

600

gttggttcat ggccattggt ctgggctctt tttgtgagtt tcatgctcgg tactgcatac

660

tctatcaatt tgccactttt acggtggaaa agatttgcat tggttgcagc aatgtgtatc

720

ctcgtgtcc gagctattat tgttcaaate gccttttatc tacatattca gacacatgtg

780

tttgggaagac caatcttggt cactaggcct cttattttcg ccactgcgtt tatgagcttt

840

ttctctgtcg ttattgcatt gtttaaggat atacctgata tcgaagggga taagatatc

900  
 ggaatccgat cattctctgt aactctgggt cagaaacggg tgttttggac atgtgttaca  
 960  
 ctacttcaaa tggcttacgc tggtgcaatt ctagttggag ccacatctcc attcatatgg  
 1020  
 agcaaagtca tctcgggttg gggcatgtt atactcgcaa caactttgtg ggctcgagct  
 1080  
 aagtcggtt atctgagtag caaaaccgaa ataacttcat gttatatgtt catatggaag  
 1140  
 ctcttttatg cagagtactt gctgttacct tttttgaagt ga  
 1182

<210> 2

<211> 393

<212> PRT

<213> Arabidopsis sp

<400> 2

Met	Glu	Ser	Leu	Leu	Ser	Ser	Ser	Ser	Leu	Val	Ser	Ala	Ala	Gly	Gly	1	5	10	15
Phe	Cys	Trp	Lys	Lys	Gln	Asn	Leu	Lys	Leu	His	Ser	Leu	Ser	Glu	Ile	20	25	30	
Arg	Val	Leu	Arg	Cys	Asp	Ser	Ser	Lys	Val	Val	Ala	Lys	Pro	Lys	Phe	35	40	45	
Arg	Asn	Asn	Leu	Val	Arg	Pro	Asp	Gly	Gln	Gly	Ser	Ser	Leu	Leu	Leu	50	55	60	
Tyr	Pro	Lys	His	Lys	Ser	Arg	Phe	Arg	Val	Asn	Ala	Thr	Ala	Gly	Gln	65	70	75	80
Pro	Glu	Ala	Phe	Asp	Ser	Asn	Ser	Lys	Gln	Lys	Ser	Phe	Arg	Asp	Ser	85	90	95	
Leu	Asp	Ala	Phe	Tyr	Arg	Phe	Ser	Arg	Pro	His	Thr	Val	Ile	Gly	Thr	100	105	110	
Val	Leu	Ser	Ile	Leu	Ser	Val	Ser	Phe	Leu	Ala	Val	Glu	Lys	Val	Ser	115	120	125	
Asp	Ile	Ser	Pro	Leu	Leu	Phe	Thr	Gly	Ile	Leu	Glu	Ala	Val	Val	Ala	130	135	140	
Ala	Leu	Met	Met	Asn	Ile	Tyr	Ile	Val	Gly	Leu	Asn	Gln	Leu	Ser	Asp	145	150	155	160
Val	Glu	Ile	Asp	Lys	Val	Asn	Lys	Pro	Tyr	Leu	Pro	Leu	Ala	Ser	Gly	165	170	175	
Glu	Tyr	Ser	Val	Asn	Thr	Gly	Ile	Ala	Ile	Val	Ala	Ser	Phe	Ser	Ile	180	185	190	
Met	Ser	Phe	Trp	Leu	Gly	Trp	Ile	Val	Gly	Ser	Trp	Pro	Leu	Phe	Trp	195	200	205	
Ala	Leu	Phe	Val	Ser	Phe	Met	Leu	Gly	Thr	Ala	Tyr	Ser	Ile	Asn	Leu	210	215	220	
Pro	Leu	Leu	Arg	Trp	Lys	Arg	Phe	Ala	Leu	Val	Ala	Ala	Met	Cys	Ile	225	230	235	240
Leu	Ala	Val	Arg	Ala	Ile	Ile	Val	Gln	Ile	Ala	Phe	Tyr	Leu	His	Ile	245	250	255	
Gln	Thr	His	Val	Phe	Gly	Arg	Pro	Ile	Leu	Phe	Thr	Arg	Pro	Leu	Ile	260	265	270	
Phe	Ala	Thr	Ala	Phe	Met	Ser	Phe	Phe	Ser	Val	Val	Ile	Ala	Leu	Phe	275	280	285	
Lys	Asp	Ile	Pro	Asp	Ile	Glu	Gly	Asp	Lys	Ile	Phe	Gly	Ile	Arg	Ser	290	295	300	
Phe	Ser	Val	Thr	Leu	Gly	Gln	Lys	Arg	Val	Phe	Trp	Thr	Cys	Val	Thr	305	310	315	320
Leu	Leu	Gln	Met	Ala	Tyr	Ala	Val	Ala	Ile	Leu	Val	Gly	Ala	Thr	Ser	325	330	335	
Pro	Phe	Ile	Trp	Ser	Lys	Val	Ile	Ser	Val	Val	Gly	His	Val	Ile	Leu				

		340					345					350			
Ala	Thr	Thr	Leu	Trp	Ala	Arg	Ala	Lys	Ser	Val	Asp	Leu	Ser	Ser	Lys
		355					360					365			
Thr	Glu	Ile	Thr	Ser	Cys	Tyr	Met	Phe	Ile	Trp	Lys	Leu	Phe	Tyr	Ala
	370					375					380				
Glu	Tyr	Leu	Leu	Leu	Pro	Phe	Leu	Lys							
385					390										

&lt;210&gt; 3

&lt;211&gt; 1224

&lt;212&gt; DNA

&lt;213&gt; Arabidopsis sp

&lt;400&gt; 3

atggcggtttt ttgggctctc ccgtgtttca agacggttgt tgaaatcttc cgtctccgta  
 60  
 actccatctt cttcctctgc tcttttgcaa tcacaacata aatccttgtc caatcctgtg  
 120  
 actaccatt acacaaatcc ttctactaag tggtatcctt catggaatga taattaccaa  
 180  
 gtatggagta aaggaagaga attgcatcag gagaagtttt ttggtgttgg ttggaattac  
 240  
 agattaattt gtggaatgtc gtcgtcttct tcggttttgg agggaaagcc gaagaaagat  
 300  
 gataaggaga agagtgatgg tggtgttgtt aagaaagctt cttggataga tttgtattta  
 360  
 ccagaagaag ttagaggtta tgctaagctt gctcgattgg ataaacccat tggaacttgg  
 420  
 ttgcttgctg ggccttgtat gtggtcgatt gcgttggtg ctgaccttgg aagccttcca  
 480  
 agttttaaat atatggcttt atttggttgc ggagcattac ttcttagagg tgctggttgt  
 540  
 actataaatg atctgcttga tcaggacata gatacaaagg ttgatcgtac aaaactaaga  
 600  
 cctatcgcca gtggtctttt gacaccattt caagggattg gatttctcgg gctgcagttg  
 660  
 ctttttaggt tagggattct tctccaactt aacaattaca gccgtgtttt aggggcttca  
 720  
 tctttgttac ttgtcttttc ctaccactt atgaagaggt ttacattttg gcctcaagcc  
 780  
 ttttttaggt tgaccataaa ctggggagca ttgttaggat ggactgcagt taaaggaagc  
 840  
 atagcaccat ctattgtact ccctctctat ctctccggag tctgctggac ccttggttat  
 900  
 gatactattt atgcacatca ggacaaagaa gatgatgtaa aagttggtgt taagtcaaca  
 960  
 gcccttagat tcggtgataa tacaaagctt tggtaactg gatttggcac agcatccata  
 1020  
 ggttttcttg cactttcttg attcagtgca gatctcgggt ggcaatatta cgcactcactg  
 1080  
 gccgctgcat caggacagtt aggatggcaa atagggacag ctgacttata atctggtgct  
 1140  
 gactgcagta gaaaatttgt gtcgaacaag tggtttgggt ctattatatt tagtggagtt  
 1200  
 gtacttgga gaagttttca ataa  
 1224

&lt;210&gt; 4

&lt;211&gt; 407

&lt;212&gt; PRT

&lt;213&gt; Arabidopsis sp

&lt;400&gt; 4

Met Ala Phe Phe Gly Leu Ser Arg Val Ser Arg Arg Leu Leu Lys Ser  
 1 5 10 15  
 Ser Val Ser Val Thr Pro Ser Ser Ser Ser Ala Leu Leu Gln Ser Gln  
 20 25 30  
 His Lys Ser Leu Ser Asn Pro Val Thr Thr His Tyr Thr Asn Pro Phe  
 35 40 45  
 Thr Lys Cys Tyr Pro Ser Trp Asn Asp Asn Tyr Gln Val Trp Ser Lys  
 50 55 60  
 Gly Arg Glu Leu His Gln Glu Lys Phe Phe Gly Val Gly Trp Asn Tyr  
 65 70 75 80  
 Arg Leu Ile Cys Gly Met Ser Ser Ser Ser Ser Val Leu Glu Gly Lys  
 85 90 95  
 Pro Lys Lys Asp Asp Lys Glu Lys Ser Asp Gly Val Val Val Lys Lys  
 100 105 110  
 Ala Ser Trp Ile Asp Leu Tyr Leu Pro Glu Glu Val Arg Gly Tyr Ala  
 115 120 125  
 Lys Leu Ala Arg Leu Asp Lys Pro Ile Gly Thr Trp Leu Leu Ala Trp  
 130 135 140  
 Pro Cys Met Trp Ser Ile Ala Leu Ala Ala Asp Pro Gly Ser Leu Pro  
 145 150 155 160  
 Ser Phe Lys Tyr Met Ala Leu Phe Gly Cys Gly Ala Leu Leu Leu Arg  
 165 170 175  
 Gly Ala Gly Cys Thr Ile Asn Asp Leu Leu Asp Gln Asp Ile Asp Thr  
 180 185 190  
 Lys Val Asp Arg Thr Lys Leu Arg Pro Ile Ala Ser Gly Leu Leu Thr  
 195 200 205  
 Pro Phe Gln Gly Ile Gly Phe Leu Gly Leu Gln Leu Leu Leu Gly Leu  
 210 215 220  
 Gly Ile Leu Leu Gln Leu Asn Asn Tyr Ser Arg Val Leu Gly Ala Ser  
 225 230 235 240  
 Ser Leu Leu Leu Val Phe Ser Tyr Pro Leu Met Lys Arg Phe Thr Phe  
 245 250 255  
 Trp Pro Gln Ala Phe Leu Gly Leu Thr Ile Asn Trp Gly Ala Leu Leu  
 260 265 270  
 Gly Trp Thr Ala Val Lys Gly Ser Ile Ala Pro Ser Ile Val Leu Pro  
 275 280 285  
 Leu Tyr Leu Ser Gly Val Cys Trp Thr Leu Val Tyr Asp Thr Ile Tyr  
 290 295 300  
 Ala His Gln Asp Lys Glu Asp Asp Val Lys Val Gly Val Lys Ser Thr  
 305 310 315 320  
 Ala Leu Arg Phe Gly Asp Asn Thr Lys Leu Trp Leu Thr Gly Phe Gly  
 325 330 335  
 Thr Ala Ser Ile Gly Phe Leu Ala Leu Ser Gly Phe Ser Ala Asp Leu  
 340 345 350  
 Gly Trp Gln Tyr Tyr Ala Ser Leu Ala Ala Ala Ser Gly Gln Leu Gly  
 355 360 365  
 Trp Gln Ile Gly Thr Ala Asp Leu Ser Ser Gly Ala Asp Cys Ser Arg  
 370 375 380  
 Lys Phe Val Ser Asn Lys Trp Phe Gly Ala Ile Ile Phe Ser Gly Val  
 385 390 395 400  
 Val Leu Gly Arg Ser Phe Gln  
 405

&lt;210&gt; 5

&lt;211&gt; 1296

&lt;212&gt; DNA

&lt;213&gt; Arabidopsis sp

&lt;400&gt; 5

atgtggcgaa gatctgttgt ttctcgttta tcttcaagaa tctctgtttc ttcttcgtta  
 60  
 ccaaacccta gactgattcc ttggtcccgc gaattatgtg ccgttaatag cttctcccag  
 120  
 cctccggtct cgacggaatc aactgctaag ttagggatca ctggtgttag atctgatgcc  
 180  
 aatcgagttt ttgccactgc tactgccgcc gctacagcta cagctaccac cggtgagatt  
 240  
 tcgtctagag ttgcggcttt ggctggatta gggcatcact acgctcgttg ttattgggag  
 300  
 ctttctaaag ctaaacttag tatgcttgtg gttgcaactt ctggaactgg gtatattctg  
 360  
 ggtacgggaa atgctgcaat tagcttcccg gggctttgtt acacatgtgc aggaaccatg  
 420  
 atgattgctg catctgctaa ttccctgaat cagatttttg agataagcaa tgattctaag  
 480  
 atgaaaagaa cgatgctaag gccattgcct tcaggacgta ttagtggtcc acacgctgtt  
 540  
 gcatgggcta ctattgctgg tgcttctggg gcttggttgg tggccagcaa gactaatatg  
 600  
 ttggctgctg gacttgcatc tgccaatctt gtactttatg cgtttgttta tactccgttg  
 660  
 aagcaacttc accctatcaa tacatgggtt ggcgctgttg ttggtgctat cccacccttg  
 720  
 cttgggtggg cggcagcgtc tggtcagatt tcatacaatt cgatgattct tccagctgct  
 780  
 ctttactttt ggcagatacc tcattttatg gcccttgac atctctgccg caatgattat  
 840  
 gcagctggag gttacaagat gttgtcactc tttgatccgt cagggaagag aatagcagca  
 900  
 gtggctctaa ggaactgctt ttacatgac cctctcgggt tcatcgcta tgactggggg  
 960  
 ttaacctcaa gttgggtttg cctcgaatca acacttctca cactagcaat cgctgcaaca  
 1020  
 gcattttcat tctaccgaga ccggaccatg cataaagcaa ggaaaatgtt ccatgccagt  
 1080  
 cttctcttcc ttctgtttt catgtctggg cttcttctac accgtgtctc taatgataat  
 1140  
 cagcaacaac tcgtagaaga agccggatta acaaattctg tatctgggtga agtcaaaact  
 1200  
 cagaggcgaa agaaacgtgt ggctcaacct ccggtggctt atgcctctgc tgcaccgttt  
 1260  
 cctttcctcc cagctccttc cttctactct ccatga  
 1296

&lt;210&gt; 6

&lt;211&gt; 431

&lt;212&gt; PRT

&lt;213&gt; Arabidopsis sp

&lt;400&gt; 6

Met	Trp	Arg	Arg	Ser	Val	Val	Tyr	Arg	Phe	Ser	Ser	Arg	Ile	Ser	Val
1				5					10					15	
Ser	Ser	Ser	Leu	Pro	Asn	Pro	Arg	Leu	Ile	Pro	Trp	Ser	Arg	Glu	Leu
			20					25					30		
Cys	Ala	Val	Asn	Ser	Phe	Ser	Gln	Pro	Pro	Val	Ser	Thr	Glu	Ser	Thr
		35					40					45			
Ala	Lys	Leu	Gly	Ile	Thr	Gly	Val	Arg	Ser	Asp	Ala	Asn	Arg	Val	Phe
	50					55					60				
Ala	Thr	Ala	Thr	Ala	Ala	Ala	Thr	Ala	Thr	Ala	Thr	Thr	Gly	Glu	Ile
65					70				75					80	



Ser Ser Arg Val Ala Ala Leu Ala Gly Leu Gly His His Tyr Ala Arg  
 85 90 95  
 Cys Tyr Trp Glu Leu Ser Lys Ala Lys Leu Ser Met Leu Val Val Ala  
 100 105 110  
 Thr Ser Gly Thr Gly Tyr Ile Leu Gly Thr Gly Asn Ala Ala Ile Ser  
 115 120 125  
 Phe Pro Gly Leu Cys Tyr Thr Cys Ala Gly Thr Met Met Ile Ala Ala  
 130 135 140  
 Ser Ala Asn Ser Leu Asn Gln Ile Phe Glu Ile Ser Asn Asp Ser Lys  
 145 150 155 160  
 Met Lys Arg Thr Met Leu Arg Pro Leu Pro Ser Gly Arg Ile Ser Val  
 165 170 175  
 Pro His Ala Val Ala Trp Ala Thr Ile Ala Gly Ala Ser Gly Ala Cys  
 180 185 190  
 Leu Leu Ala Ser Lys Thr Asn Met Leu Ala Ala Gly Leu Ala Ser Ala  
 195 200 205  
 Asn Leu Val Leu Tyr Ala Phe Val Tyr Thr Pro Leu Lys Gln Leu His  
 210 215 220  
 Pro Ile Asn Thr Trp Val Gly Ala Val Val Gly Ala Ile Pro Pro Leu  
 225 230 235 240  
 Leu Gly Trp Ala Ala Ala Ser Gly Gln Ile Ser Tyr Asn Ser Met Ile  
 245 250 255  
 Leu Pro Ala Ala Leu Tyr Phe Trp Gln Ile Pro His Phe Met Ala Leu  
 260 265 270  
 Ala His Leu Cys Arg Asn Asp Tyr Ala Ala Gly Gly Tyr Lys Met Leu  
 275 280 285  
 Ser Leu Phe Asp Pro Ser Gly Lys Arg Ile Ala Ala Val Ala Leu Arg  
 290 295 300  
 Asn Cys Phe Tyr Met Ile Pro Leu Gly Phe Ile Ala Tyr Asp Trp Gly  
 305 310 315 320  
 Leu Thr Ser Ser Trp Phe Cys Leu Glu Ser Thr Leu Leu Thr Leu Ala  
 325 330 335  
 Ile Ala Ala Thr Ala Phe Ser Phe Tyr Arg Asp Arg Thr Met His Lys  
 340 345 350  
 Ala Arg Lys Met Phe His Ala Ser Leu Leu Phe Leu Pro Val Phe Met  
 355 360 365  
 Ser Gly Leu Leu Leu His Arg Val Ser Asn Asp Asn Gln Gln Gln Leu  
 370 375 380  
 Val Glu Glu Ala Gly Leu Thr Asn Ser Val Ser Gly Glu Val Lys Thr  
 385 390 395 400  
 Gln Arg Arg Lys Lys Arg Val Ala Gln Pro Pro Val Ala Tyr Ala Ser  
 405 410 415  
 Ala Ala Pro Phe Pro Phe Leu Pro Ala Pro Ser Phe Tyr Ser Pro  
 420 425 430

&lt;210&gt; 7

&lt;211&gt; 479

&lt;212&gt; DNA

&lt;213&gt; Arabidopsis sp

&lt;400&gt; 7

ggaaactccc ggagcacctg tttgcaggta ccgctaacct taatcgataa tttatttctc  
 60  
 ttgtcaggaa ttatgtaagt ctggtggaag gctcgcatat catttttgca ttgcctttcg  
 120  
 ctatgatcgg gtttactttg ggtgtgatga gaccaggcgt ggctttatgg tatggcgaaa  
 180  
 acccattttt atccaatgct gcattccctc ccgatgatcc gttctttcat tcctatacag  
 240  
 gtatcatgct gataaaactg ttactgggtac tggtttgbat ggtatcagca agaagcgcg  
 300

cgatggcggtt taaccgggat ctgcacaggc attttgacgc gaagaacccg cgtactgcca  
360  
tccgtgaaat acctgcgggc gtcatactctg ccaacagtgc gctgggtgttt acgataggct  
420  
gctgcgtggg attctgggtg gcctgttatt tcattaacac gatctgtttt tacctggcg  
479

<210> 8  
<211> 551  
<212> DNA  
<213> Arabidopsis sp

<220>  
<221> misc\_feature  
<222> (1)...(551)  
<223> n = A,T,C or G

<400> 8  
ttgtggctta caccttaatg agcatagcc agnccattac ggctcgtaa tcggcgccat  
60  
ngccgngct gntgcaccgg tagtgggcta ctgcgccgtg accaatcagc ttgatctagc  
120  
ggctcttatt ctgtttttaa ttttactgtt ctggcaaagt ccgcattttt acgcgatttc  
180  
cattttcagg ctaaaagact tttcagcggc ctgtattccg gtgctgcca tcattaaaga  
240  
cctgcgctat accaaaatca gcctgctggg ttacgtgggc ttatttacac tggctgctat  
300  
catgccggcc ctcttagggg atgccgggtg gatttatggg atagcggcct taattttagg  
360  
cttgatttgg ctttatattg ccatacaagg attcaagacc gccgatgatc aaaaatggtc  
420  
tcgtaagatg tttggatctt cgattttaat cattaccctc ttgtcggtaa tgatgcttgt  
480  
ttaaacttac tgcctcctga agtttatata tcgataattt cagcttaagg aggcttagtg  
540  
gttaattcaa t  
551

<210> 9  
<211> 297  
<212> PRT  
<213> Arabidopsis sp

<400> 9  
Met Val Leu Ala Glu Val Pro Lys Leu Ala Ser Ala Ala Glu Tyr Phe  
1 5 10 15  
Phe Lys Arg Gly Val Gln Gly Lys Gln Phe Arg Ser Thr Ile Leu Leu  
20 25 30  
Leu Met Ala Thr Ala Leu Asn Val Arg Val Pro Glu Ala Leu Ile Gly  
35 40 45  
Glu Ser Thr Asp Ile Val Thr Ser Glu Leu Arg Val Arg Gln Arg Gly  
50 55 60  
Ile Ala Glu Ile Thr Glu Met Ile His Val Ala Ser Leu Leu His Asp  
65 70 75 80  
Asp Val Leu Asp Asp Ala Asp Thr Arg Arg Gly Val Gly Ser Leu Asn  
85 90 95  
Val Val Met Gly Asn Lys Val Val Ala Leu Leu Ala Thr Ala Val Glu  
100 105 110  
His Leu Val Thr Gly Glu Thr Met Glu Ile Thr Ser Ser Thr Glu Gln  
115 120 125

Arg	Tyr	Ser	Met	Asp	Tyr	Tyr	Met	Gln	Lys	Thr	Tyr	Tyr	Lys	Thr	Ala
130						135					140				
Ser	Leu	Ile	Ser	Asn	Ser	Cys	Lys	Ala	Val	Ala	Val	Leu	Thr	Gly	Gln
145					150					155					160
Thr	Ala	Glu	Val	Ala	Val	Leu	Ala	Phe	Glu	Tyr	Gly	Arg	Asn	Leu	Gly
				165					170					175	
Leu	Ala	Phe	Gln	Leu	Ile	Asp	Asp	Ile	Leu	Asp	Phe	Thr	Gly	Thr	Ser
			180					185					190		
Ala	Ser	Leu	Gly	Lys	Gly	Ser	Leu	Ser	Asp	Ile	Arg	His	Gly	Val	Ile
		195					200					205			
Thr	Ala	Pro	Ile	Leu	Phe	Ala	Met	Glu	Glu	Phe	Pro	Gln	Leu	Arg	Glu
	210					215					220				
Val	Val	Asp	Gln	Val	Glu	Lys	Asp	Pro	Arg	Asn	Val	Asp	Ile	Ala	Leu
225					230					235					240
Glu	Tyr	Leu	Gly	Lys	Ser	Lys	Gly	Ile	Gln	Arg	Ala	Arg	Glu	Leu	Ala
				245					250					255	
Met	Glu	His	Ala	Asn	Leu	Ala	Ala	Ala	Ala	Ile	Gly	Ser	Leu	Pro	Glu
			260				265						270		
Thr	Asp	Asn	Glu	Asp	Val	Lys	Arg	Ser	Arg	Arg	Ala	Leu	Ile	Asp	Leu
		275					280					285			
Thr	His	Arg	Val	Ile	Thr	Arg	Asn	Lys							
	290					295									

&lt;210&gt; 10

&lt;211&gt; 561

&lt;212&gt; DNA

&lt;213&gt; Arabidopsis sp.

&lt;400&gt; 10

aagcgcatcc gtcctcttct acgattgccg ccagccgcat gtatggctgc ataaccgacc  
60  
gccctatcc gctcgcggcc gcggtcgaat tcattcacac cgcgacgctg ctgcatgacg  
120  
acgtcgtcga tgaaagcgat ttgcgccgcg gccgcgaaag cgcgcataag gttttcggca  
180  
atcaggcgag cgtgctcgtc ggcgatttcc ttttctccc cgccttccag ctgatgggtg  
240  
aagacggctc gctcgacgcg ctgcgcattc tctcggtatgc ctccgccgtg atcgcgcagg  
300  
gcgaagtgat gcagctcggc accgcgcgca atcttgaaac caatatgagc cagtatctcg  
360  
atgtgatcag cgcgaagacc gccgcgctct ttgccgccgc ctgcgaaatc ggcccgggtga  
420  
tggcgaacgc gaaggcgga gatgctgccg cgatgtgcga atacggcatg aatctcggta  
480  
tcgccttcca gatcatcgac gaccttctcg attacggcac cggcggccac gccgagcttg  
540  
gcaagaacac gggcgacgat t  
561

&lt;210&gt; 11

&lt;211&gt; 966

&lt;212&gt; DNA

&lt;213&gt; Arabidopsis sp

&lt;400&gt; 11

atggtacttg ccgaggttcc aaagcttgcc tctgctgctg agtacttctt caaaaggggt  
60  
gtgcaaggaa aacagtttctg ttcaactatt ttgctgctga tggcgacagc tctgaatgta  
120  
cgcgttccag aagcattgat tggggaatca acagatatag tcacatcaga attacgcgta

180  
 aggcaacggg gtattgctga aatcactgaa atgatacacg tgcgaagtct actgcacgat  
 240  
 gatgtcttgg atgatgccga tacaaggcgt ggtgttggtt ccttaaagtgt tgtaatgggt  
 300  
 aacaagatgt cggatttagc aggagacttc ttgctctccc gggcttgtgg ggctctcgct  
 360  
 gcttttaaaga acacagaggt tgtagcatta cttgcaactg ctgtagaaca tcttgttacc  
 420  
 ggtgaaacca tggaaataac tagttcaacc gagcagcgtt atagtatgga ctactacatg  
 480  
 cagaagacat attataagac agcatcgcta atctctaaca gctgcaaagc tggtgccgtt  
 540  
 ctactggac aaacagcaga agttgccgtg ttagcttttg agtatgggag gaatctgggt  
 600  
 ttagcattcc aattaataga cgacattctt gatttcacgg gcacatctgc ctctctcgga  
 660  
 aagggatcgt tgtcagatat tgcctatgga gtcataacag cccaatcct ctttgccatg  
 720  
 gaagagtttc ctcaactacg cgaagttggt gatcaagttg aaaaagatcc taggaatggt  
 780  
 gacattgctt tagagtatct tgggaagagc aagggaatac agagggcaag agaattagcc  
 840  
 atggaacatg cgaatctagc agcagctgca atcgggtctc tacctgaaac agacaatgaa  
 900  
 gatgtcaaaa gatcgaggcg ggcacttatt gacttgaccc atagagtcac caccagaaac  
 960  
 aagtga  
 966

<210> 12  
 <211> 321  
 <212> PRT  
 <213> Arabidopsis sp

<400> 12  
 Met Val Leu Ala Glu Val Pro Lys Leu Ala Ser Ala Ala Glu Tyr Phe  
 1 5 10 15  
 Phe Lys Arg Gly Val Gln Gly Lys Gln Phe Arg Ser Thr Ile Leu Leu  
 20 25 30  
 Leu Met Ala Thr Ala Leu Asn Val Arg Val Pro Glu Ala Leu Ile Gly  
 35 40 45  
 Glu Ser Thr Asp Ile Val Thr Ser Glu Leu Arg Val Arg Gln Arg Gly  
 50 55 60  
 Ile Ala Glu Ile Thr Glu Met Ile His Val Ala Ser Leu Leu His Asp  
 65 70 75 80  
 Asp Val Leu Asp Asp Ala Asp Thr Arg Arg Gly Val Gly Ser Leu Asn  
 85 90 95  
 Val Val Met Gly Asn Lys Met Ser Val Leu Ala Gly Asp Phe Leu Leu  
 100 105 110  
 Ser Arg Ala Cys Gly Ala Leu Ala Ala Leu Lys Asn Thr Glu Val Val  
 115 120 125  
 Ala Leu Leu Ala Thr Ala Val Glu His Leu Val Thr Gly Glu Thr Met  
 130 135 140  
 Glu Ile Thr Ser Ser Thr Glu Gln Arg Tyr Ser Met Asp Tyr Tyr Met  
 145 150 155 160  
 Gln Lys Thr Tyr Tyr Lys Thr Ala Ser Leu Ile Ser Asn Ser Cys Lys  
 165 170 175  
 Ala Val Ala Val Leu Thr Gly Gln Thr Ala Glu Val Ala Val Leu Ala  
 180 185 190  
 Phe Glu Tyr Gly Arg Asn Leu Gly Leu Ala Phe Gln Leu Ile Asp Asp

195	200	205
Ile Leu Asp Phe Thr Gly Thr Ser Ala Ser Leu Gly Lys Gly Ser Leu		
210	215	220
Ser Asp Ile Arg His Gly Val Ile Thr Ala Pro Ile Leu Phe Ala Met		
225	230	235
Glu Glu Phe Pro Gln Leu Arg Glu Val Val Asp Gln Val Glu Lys Asp		
245	250	255
Pro Arg Asn Val Asp Ile Ala Leu Glu Tyr Leu Gly Lys Ser Lys Gly		
260	265	270
Ile Gln Arg Ala Arg Glu Leu Ala Met Glu His Ala Asn Leu Ala Ala		
275	280	285
Ala Ala Ile Gly Ser Leu Pro Glu Thr Asp Asn Glu Asp Val Lys Arg		
290	295	300
Ser Arg Arg Ala Leu Ile Asp Leu Thr His Arg Val Ile Thr Arg Asn		
305	310	315
Lys		320

<210> 13  
 <211> 621  
 <212> DNA  
 <213> Arabidopsis sp

<400> 13  
 gctttctcct ttgctaattc ttgagctttc ttgatccac cgcgatttct aactatttca  
 60  
 atcgcttctt caagcgatcc aggctcacia aactcagact caatgatctc tcttagcctt  
 120  
 ggctcattct ctagcgcgaa gatcactggc gccgttatgt tacctttggc taagtcatta  
 180  
 gctgcaggct tacctaactg ctctgtggac tgagtgaagt ccagaatgtc atcaactact  
 240  
 tgaaaagata aaccgagatt cttcccgaac tgatacattt gctctgagac cttgctttcg  
 300  
 actttactga aaattgctgc tcctttggtg cttgcagcta ctaatgaagc tgtcttgtag  
 360  
 taactcttta gcatgtagtc atcaagcttg acatcacaat cgaataaact cgatgcttgc  
 420  
 tttatctcac cgcttgcaaa atctttgatc acctgcaaaa agataaatca agattcagac  
 480  
 caaatgttct ttgtattgag tagcttcac taatctcaga aaggaatatt acctgactta  
 540  
 tgagcttaat gacttcaagg ttttcgagat ttgtaagtac catgatgctt gagcaacatg  
 600  
 aaatccccag ctaatacagc t  
 621

<210> 14  
 <211> 741  
 <212> DNA  
 <213> Arabidopsis sp

<400> 14  
 ggtgagtttt gttaatagtt atgagattca tctatttttg tcataaaatt gtttggtttg  
 60  
 gtttaaactc tgtgtataat tgcaggaaag gaaacagttc atgagctttt cggcacaaga  
 120  
 gtagcgggtgc tagctggaga tttcatgttt gctcaagcgt catggtactt agcaaacttc  
 180  
 gagaatcttg aagttattaa gctcatcagt caggactta gttactctta cattgttttt  
 240

ctatgagggtt gagctatgaa tctcatttcg ttgaataatg ctgtgcctca aacttttttt  
300  
catgttttca ggtgatcaaa gactttgcaa gcggagagat aaagcaggcg tccagcttat  
360  
ttgactgcga caccaagctc gacgagtact tactcaaaag tttctacaag acagcctctt  
420  
tagtggctgc gagcaccaaa ggagctgcc a ttttcagcag agttgagcct gatgtgacag  
480  
aacaatgta cgagtttggg aagaatctcg gtctctcttt ccagatagtt gatgatattt  
540  
tggatttcac tcagtcgaca gacgagctcg ggaagccagc agggagtgat ttggctaaag  
600  
gtaacttaac agcacctgtg attttcgctc tggagaggga gccaaggcta agagagatca  
660  
ttgagtcaaa gttctgtgag gcgggttctc tggaagaagc gattgaagcg gtgacaaaag  
720  
gtggggggat taagagagca c  
741

&lt;210&gt; 15

&lt;211&gt; 1087

&lt;212&gt; DNA

&lt;213&gt; Arabidopsis sp

&lt;400&gt; 15

cctcttcagc caatccagag gaagaagaga caacttttta tctttcgtca agagtctccg  
60  
aaaacgcacg gttttatgct ctctcttctg ccctcacctc acaagacgca gggcacatga  
120  
ttcaaccaga gggaaaaagc aacgataaca actctgcttt tgatttcaag ctgtatatga  
180  
tccgcaaagc cgagtctgta aatgcggctc tcgacgtttc cgtaccgctt ctgaaacccc  
240  
ttacgatcca agaagcggtc aggtactctt tgctagccgg cggaaaacgt gtgaggcctc  
300  
tgctctgcat tgccgcttgt gagcttgtgg ggggcgacga ggctactgcc atgtcagccg  
360  
cttgcgcggt cgagatgatc cacacaagct ctctcattca tgacgatctt ccgtgcatgg  
420  
acaatgccga cctccgtaga ggcaagccca ccaatcacia ggtatgttgt ttaattatat  
480  
gaaggctcag agataatgct gaactagtgt tgaaccaatt tttgctcaaa caaggatat  
540  
ggagaagaca tggcggtttt ggcagggtgat gcactccttg cattggcggt tgagcacatg  
600  
acggttgtgt cgagtgggtt ggtcgtccc gagaagatga ttcgcgccgt ggttgagctg  
660  
gccagggcca tagggactac agggctagtt gctggacaaa tgatagacct agccagcgaa  
720  
agactgaatc cagacaaggt tggattggag catctagagt tcatccatct ccacaaaacg  
780  
gcggcattgt tggaggcagc ggcagtttta ggggttataa tgggaggtgg aacagaggaa  
840  
gaaatcgaaa agcttagaaa gtatgctagg tgtattggac tactgtttca ggttggtgat  
900  
gacattctcg acgtaacaaa atctactgag gaattgggta agacagccgg aaaagacgta  
960  
atggccggaa agctgacgta tccaaggctg ataggtttgg agggatccag ggaagttgca  
1020  
gagcacctga ggagagaagc agaggaaaag cttaaagggt ttgatccaag tcaggcgggc  
1080



cctctgg  
1087

<210> 16  
<211> 1164  
<212> DNA  
<213> Arabidopsis sp

<400> 16  
atgacttcga ttctcaacac tgtctccacc atccactctt ccagagttac ctccgtcgat  
60  
cgagtcggag tcctctctct tcggaattcg gattccgttg agttcactcg ccggcggttct  
120  
ggtttctcga cgttgatcta cgaatcaccg gggcggagat ttgttggtgcg tgcggcggag  
180  
actgatactg ataaagttaa atctcagaca cctgacaagg caccagccgg tgggtcaagc  
240  
attaaccagc ttctcggtat caaaggagca tctcaagaaa ctaataaatg gaagattcgt  
300  
cttcagctta caaaaccagt cacttggcct ccactggttt ggggagtcgt ctgtggtgct  
360  
gctgcttcag ggaactttca ttggacccca gaggatgttg ctaagtcgat tctttgcatg  
420  
atgatgtctg gtccttgtct tactggctat acacagacaa tcaacgactg gtatgataga  
480  
gatatcgacg caattaatga gccatategt ccaattccat ctggagcaat atcagagcca  
540  
gaggttatta cacaagtctg ggtgctatta ttgggaggtc ttggtattgc tggaatatta  
600  
gatgtgtggg cagggcatac cactcccact gtcttctatc ttgctttggg aggatcattg  
660  
ctatcttata tatactctgc tccacctctt aagctaaaac aaaatggatg gggttgaaat  
720  
tttgacttg gagcaagcta tattagtttg ccatgggtggg ctggccaagc attgtttggc  
780  
actcttacgc cagatgttgt tgttctaaca ctcttgtaca gcatagctgg gttaggaata  
840  
gccattgtta acgacttcaa aagtgttgaa ggagatagag cattaggact tcagtctctc  
900  
ccagtagctt ttggcaccga aactgcaaaa tggatatgcg ttggtgctat agacattact  
960  
cagctttctg ttgccggata tctattagca tctgggaaac cttattatgc gttggcggtg  
1020  
gttgctttga tcattcctca gattgtgttc cagtttaaata actttctcaa ggaccctgtc  
1080  
aaatacgacg tcaagtacca ggcaagcgcg cagccattct tgggtgctcg aatatttgta  
1140  
acggcattag catcgcaaca ctga  
1164

<210> 17  
<211> 387  
<212> PRT  
<213> Arabidopsis sp

<400> 17  
Met Thr Ser Ile Leu Asn Thr Val Ser Thr Ile His Ser Ser Arg Val  
1 5 10 15  
Thr Ser Val Asp Arg Val Gly Val Leu Ser Leu Arg Asn Ser Asp Ser  
20 25 30  
Val Glu Phe Thr Arg Arg Arg Ser Gly Phe Ser Thr Leu Ile Tyr Glu

```
<400> 18
atggttgttta gtggttcagc gatcccatta agcagcttct gctctcttcc ggagaaaccc
60
cacactcttc ctatgaaact ctctcccgtc gcaatccgat cttcatcctc atctgccccg
120
gggtcgttga acttcgatct gaggacgtat tggacgactc tgatcaccca gatcaaccag
180
aagctggatg aggccatacc ggtcaagcac cctgcgggga tctacgaggc tatgagatac
240
tctgtactcg cacaaggcgc caagcgtgcc cctcctgtga tgtgtgtggc ggcctgcgag
```

300  
ctcttcggtg gcgatcgctt cgccgctttc cccaccgcct gtgccctaga aatggtgcac  
360  
gcggcttcgt tgatacacga cgacctcccc tgtatggacg acgatcctgt gcgcagagga  
420  
aagccatcta accacactgt ctacggctct ggcatggcca ttctcgccgg tgacgccctc  
480  
ttcccactcg ccttcacga cattgtctcc cacacgcctc ctgaccttgt tccccgagcc  
540  
accatcctca gactcatcac tgagattgcc cgcactgtcg gctccactgg tatggctgca  
600  
ggccagtacg tcgaccttga aggaggctccc ttctctcttt cctttgttca ggagaagaaa  
660  
ttcggagcca tgggtgaatg ctctgccgtg tgcggtggcc tattgggagg tgccactgag  
720  
gatgagctcc agagtctccg aaggtaaggg agagccgtcg ggatgctgta tcaggtggtc  
780  
gatgacatca ccgaggacaa gaagaagagc tatgatgggtg gagcagagaa gggaatgatg  
840  
gaaatggcgg aagagctcaa ggagaaggcg aagaaggagc ttcaagtgtt tgacaacaag  
900  
tatggaggag gagacacact tgttctcttc tacaccttcg ttgactacgc tgctcatcga  
960  
cattttcttc tccccctctg a  
981

<210> 19  
<211> 245  
<212> DNA  
<213> GLycine sp

<400> 19  
gcaacatctg ggactgggtt tgtcttgggg agtggtagtg ctgttgatct ttcggcactt  
60  
tcttgacttt gcttgggtac catgatggtt gctgcatctg ctaactcttt gaatcaggtg  
120  
tttgagatca ataatgatgc taaaatgaag agaacaagtc gcaggccact accctcagga  
180  
cgcatacaaa tacctcatgc agttggctgg gcatactctg ttggattagc tggtagggct  
240  
ctact  
245

<210> 20  
<211> 253  
<212> DNA  
<213> Glycine sp

<400> 20  
attggctttc caagatcatt gggttttctt gttgcattca tgaccttcta ctcttgggt  
60  
ttggcattgt ccaaggatat acctgacgtt gaaggagata aagagcacgg cattgattct  
120  
tttgagtagc gtctaggtca gaaacgggca ttttggattt gcgtttcctt ttttgaaatg  
180  
gctttcggag ttggtatcct ggccggagca tcatgctcac acttttggac taaaattttc  
240  
acgggtatgg gaa  
253

<210> 21

<211> 275  
<212> DNA  
<213> Glycine sp

<400> 21  
tgatcttcta ctctctgggt atggcattgt ccaaggatat atctgacgtt aaaggagata  
60  
aagcatacgg catcgatact ttagcgatac gtttgggtca aaaatgggta ttttggattt  
120  
gcattatcct ttttgaaatg gcttttggag ttgccctctt ggcaggagca acatcttctt  
180  
acctttggat taaaattgtc acgggtctgg gacatgctat tcttgcttca attctcttgt  
240  
accaagccaa atctatatac ttgagcaaca aagtt  
275

<210> 22  
<211> 299  
<212> DNA  
<213> Glycine sp

<220>  
<221> misc feature  
<222> (1) ... (299)  
<223> n = A,T,C or G

<400> 22  
ccanaatang tncatcttng aaagacaatt ggcctcttca acacacaagt ctgcatgtga  
60  
agaagaggcc aattgtcttt ccaagatcac ttatngtggc tattgtaatc atgaacttct  
120  
tctttgtggg tatggcattg gcaaaggata tacctanctg ttgaaggaga taaaatatat  
180  
ggcattgata cttttgcaat acgtataggt caaaaacaag tatttttggat ttgtattttc  
240  
ctttttgaaa ggctttcgga gtttccctag tggcaggagc aacatcttct agccttggg  
299

<210> 23  
<211> 767  
<212> DNA  
<213> Glycine sp

<400> 23  
gtggaggctg tggttgctgc cctgtttatg aatatttata ttgttggttt gaatcaattg  
60  
tctgatgttg aaatagacaa gataaacaag ccgtatcttc cattagcatc tggggaatat  
120  
tcctttgaaa ctggtgtcac tattgttgca tctttttcaa ttctgagttt ttggcttggc  
180  
tgggtttag gttcatggcc attatttttg gccctttttg taagctttgt gctaggaact  
240  
gcttattcaa tcaatgtgcc tctgttgaga tggagagggt ttgcagtgtc tgcagcgatg  
300  
tgcattctag ctgttcgggc agtaatagtt caacttgcac ttttccttca catgcagact  
360  
catgtgtaca agaggccacc tgtcttttca agaccattga tttttgctac tgcattcatg  
420  
agcttcttct ctgtagttaa agcactgttt aaggatatac ctgacattga aggagataaa  
480  
gtatttggca tccaatcttt ttcagtgtgt ttaggtcaga agccggtgtt ctggacttgt

540  
 gttacccttc ttgaaatagc ttatggagtc gccctcctgg tgggagctgc atctccttgt  
 600  
 ctttggagca aaattttcac gggctctggga cacgctgtgc tggcttcaat tctctgggtt  
 660  
 catgccaaat ctgtagattt gaaaagcaaa gcttcgataa catccttcta tatgtttatt  
 720  
 tggaagctat tttatgcaga atacttactc attccttttg ttagatg  
 767

<210> 24  
 <211> 255  
 <212> PRT  
 <213> Glycine sp

<400> 24  
 Val Glu Ala Val Val Ala Ala Leu Phe Met Asn Ile Tyr Ile Val Gly  
 1 5 10 15  
 Leu Asn Gln Leu Ser Asp Val Glu Ile Asp Lys Ile Asn Lys Pro Tyr  
 20 25 30  
 Leu Pro Leu Ala Ser Gly Glu Tyr Ser Phe Glu Thr Gly Val Thr Ile  
 35 40 45  
 Val Ala Ser Phe Ser Ile Leu Ser Phe Trp Leu Gly Trp Val Val Gly  
 50 55 60  
 Ser Trp Pro Leu Phe Trp Ala Leu Phe Val Ser Phe Val Leu Gly Thr  
 65 70 75 80  
 Ala Tyr Ser Ile Asn Val Pro Leu Leu Arg Trp Lys Arg Phe Ala Val  
 85 90 95  
 Leu Ala Ala Met Cys Ile Leu Ala Val Arg Ala Val Ile Val Gln Leu  
 100 105 110  
 Ala Phe Phe Leu His Met Gln Thr His Val Tyr Lys Arg Pro Pro Val  
 115 120 125  
 Phe Ser Arg Pro Leu Ile Phe Ala Thr Ala Phe Met Ser Phe Phe Ser  
 130 135 140  
 Val Val Ile Ala Leu Phe Lys Asp Ile Pro Asp Ile Glu Gly Asp Lys  
 145 150 155 160  
 Val Phe Gly Ile Gln Ser Phe Ser Val Cys Leu Gly Gln Lys Pro Val  
 165 170 175  
 Phe Trp Thr Cys Val Thr Leu Leu Glu Ile Ala Tyr Gly Val Ala Leu  
 180 185 190  
 Leu Val Gly Ala Ala Ser Pro Cys Leu Trp Ser Lys Ile Phe Thr Gly  
 195 200 205  
 Leu Gly His Ala Val Leu Ala Ser Ile Leu Trp Phe His Ala Lys Ser  
 210 215 220  
 Val Asp Leu Lys Ser Lys Ala Ser Ile Thr Ser Phe Tyr Met Phe Ile  
 225 230 235 240  
 Trp Lys Leu Phe Tyr Ala Glu Tyr Leu Leu Ile Pro Phe Val Arg  
 245 250 255

<210> 25  
 <211> 360  
 <212> DNA  
 <213> Zea sp

<220>  
 <221> misc\_feature  
 <222> (1)...(360)  
 <223> n = A,T,C or G

<400> 25  
 ggcgtcttca cttgttctgg tcttctcgta tcccctgatg aagaggttca cattttggcc

60  
 tcaggcttat cttggcctga cattcaactg gggagcttta ctaggggtggg ctgctattaa  
 120  
 ggaaagcata gaccctgcaa atcatccttc cattgtatac agctgggtatt tgttggacgc  
 180  
 tgggtgatga tactatatat gcgcatacagg tgtttcgcta tccctacttt catattaatc  
 240  
 cttgatgaag tggccatttc atgttgtcgc ggtgggtctta tacttgcata tctccatgca  
 300  
 tctcaggaca aagangatga cctgaaagta ggagtccaag tccacagctt aagatttggg  
 360

<210> 26  
 <211> 299  
 <212> DNA  
 <213> Zea sp

<220>  
 <221> misc\_feature  
 <222> (1)...(299)  
 <223> n = A,T,C or G

<400> 26  
 gatggttgca gcatctgcaa ataccctcaa ccagggtgtt gngataaaaa atgatgctaa  
 60  
 aatgaaaagg acaatgcgtg cccctgccca tctggctgca ttagtcctgc acatgctgcg  
 120  
 atgtgggcta caagtgttgg agttgcagga acagctttgt tggcctggaa ggctaattggc  
 180  
 ttggcagctg ggcttgcagc ttctaactct gttctgtatg catttgtgta tacgccgttg  
 240  
 aagcaaatac accctgttaa tacatgggtt ggggcagtcg ttggtgccat cccaccact  
 299

<210> 27  
 <211> 255  
 <212> DNA  
 <213> Zea sp

<220>  
 <221> misc\_feature  
 <222> (1)...(255)  
 <223> n = A,T,C or G

<400> 27  
 anacttgcat atctccatgc ntctcaggac aaagangatg acctgaaagt aggtgtcaag  
 60  
 tccacagcat taagatttgg agatttgacc nnatactgna tcagtggctt tggcgcgga  
 120  
 tgcttcggca gcttagcact cagtgggttac aatgctgacc ttggttggtg tttagtgtga  
 180  
 tgcttgagcg aagaatggta tngtttttac ttgatattga ctccagacct gaaatcatgt  
 240  
 tggacagggt ggccc  
 255

<210> 28  
 <211> 257  
 <212> DNA  
 <213> Zea sp



&lt;400&gt; 28

attgaagggg ataggactct ggggcttcag tcacttcctg ttgcttttgg gatggaaact  
60  
gcaaaatgga tttgtgttgg agcaattgat atcactcaat tatctgttgc aggttaccta  
120  
ttgagcaccg gtaagctgta ttatgccctg gtgttgcttg ggctaacaat tcctcaggtg  
180  
ttctttcagt tccagtactt cctgaaggac cctgtgaagt atgatgtcaa atatcaggca  
240  
agcgacaac cattctt  
257

&lt;210&gt; 29

&lt;211&gt; 368

&lt;212&gt; DNA

&lt;213&gt; Zea sp

&lt;400&gt; 29

atccagttgc aaataataat ggcgttcttc tctgttgtaa tagcactatt caaggatata  
60  
cctgacatcg aaggggaccg catattcggg atccgatact tcagcgtccg gttagggcaa  
120  
aagaaggtct tttggatctg cgttggcttg cttgagatgg cctacagcgt tgcgatactg  
180  
atgggagcta cctcttcctg tttgtggagc aaaacagcaa ccatcgctgg ccattccata  
240  
cttgccgcga tcctatggag ctgcgcgcga tcggtggact tgacgagcaa agccgcaata  
300  
acgtccttct acatgttcat ctggaagctg ttctacgcgg agtacctgct catccctctg  
360  
gtgcggtg  
368

&lt;210&gt; 30

&lt;211&gt; 122

&lt;212&gt; PRT

&lt;213&gt; Zea sp

&lt;400&gt; 30

Ile	Gln	Leu	Gln	Ile	Ile	Met	Ala	Phe	Phe	Ser	Val	Val	Ile	Ala	Leu
1			5					10					15		
Phe	Lys	Asp	Ile	Pro	Asp	Ile	Glu	Gly	Asp	Arg	Ile	Phe	Gly	Ile	Arg
		20					25					30			
Ser	Phe	Ser	Val	Arg	Leu	Gly	Gln	Lys	Lys	Val	Phe	Trp	Ile	Cys	Val
		35				40					45				
Gly	Leu	Leu	Glu	Met	Ala	Tyr	Ser	Val	Ala	Ile	Leu	Met	Gly	Ala	Thr
	50				55					60					
Ser	Ser	Cys	Leu	Trp	Ser	Lys	Thr	Ala	Thr	Ile	Ala	Gly	His	Ser	Ile
65				70				75						80	
Leu	Ala	Ala	Ile	Leu	Trp	Ser	Cys	Ala	Arg	Ser	Val	Asp	Leu	Thr	Ser
			85					90					95		
Lys	Ala	Ala	Ile	Thr	Ser	Phe	Tyr	Met	Phe	Ile	Trp	Lys	Leu	Phe	Tyr
		100					105						110		
Ala	Glu	Tyr	Leu	Leu	Ile	Pro	Leu	Val	Arg						
		115					120								

&lt;210&gt; 31

&lt;211&gt; 278

&lt;212&gt; DNA

&lt;213&gt; Zea sp

&lt;400&gt; 31

tattcagcac cacctctcaa gctcaagcag aatggatgga ttgggaactt cgctctgggt  
60  
gcgagttaca tcagcttgcc ctgggtgggct ggccaggcgt tatttggaac tcttacacca  
120  
gatatacattg tcttgactac tttgtacagc atagctgggc tagggattgc tattgtaaat  
180  
gatttcaaga gtattgaagg ggataggact ctggggcttc agtcacttcc tgttgctttt  
240  
gggatggaaa ctgcaaaatg gatttgtgtt ggagcaat  
278

&lt;210&gt; 32

&lt;211&gt; 292

&lt;212&gt; PRT

&lt;213&gt; Synechocystis sp

&lt;400&gt; 32

Met	Val	Ala	Gln	Thr	Pro	Ser	Ser	Pro	Pro	Leu	Trp	Leu	Thr	Ile	Ile
1				5					10					15	
Tyr	Leu	Leu	Arg	Trp	His	Lys	Pro	Ala	Gly	Arg	Leu	Ile	Leu	Met	Ile
			20					25					30		
Pro	Ala	Leu	Trp	Ala	Val	Cys	Leu	Ala	Ala	Gln	Gly	Leu	Pro	Pro	Leu
		35					40					45			
Pro	Leu	Leu	Gly	Thr	Ile	Ala	Leu	Gly	Thr	Leu	Ala	Thr	Ser	Gly	Leu
	50					55					60				
Gly	Cys	Val	Val	Asn	Asp	Leu	Trp	Asp	Arg	Asp	Ile	Asp	Pro	Gln	Val
65					70				75						80
Glu	Arg	Thr	Lys	Gln	Arg	Pro	Leu	Ala	Ala	Arg	Ala	Leu	Ser	Val	Gln
				85					90					95	
Val	Gly	Ile	Gly	Val	Ala	Leu	Val	Ala	Leu	Leu	Cys	Ala	Ala	Gly	Leu
			100					105					110		
Ala	Phe	Tyr	Leu	Thr	Pro	Leu	Ser	Phe	Trp	Leu	Cys	Val	Ala	Ala	Val
		115					120					125			
Pro	Val	Ile	Val	Ala	Tyr	Pro	Gly	Ala	Lys	Arg	Val	Phe	Pro	Val	Pro
	130					135					140				
Gln	Leu	Val	Leu	Ser	Ile	Ala	Trp	Gly	Phe	Ala	Val	Leu	Ile	Ser	Trp
145					150				155						160
Ser	Ala	Val	Thr	Gly	Asp	Leu	Thr	Asp	Ala	Thr	Trp	Val	Leu	Trp	Gly
				165					170					175	
Ala	Thr	Val	Phe	Trp	Thr	Leu	Gly	Phe	Asp	Thr	Val	Tyr	Ala	Met	Ala
			180					185					190		
Asp	Arg	Glu	Asp	Asp	Arg	Arg	Ile	Gly	Val	Asn	Ser	Ser	Ala	Leu	Phe
		195					200					205			
Phe	Gly	Gln	Tyr	Val	Gly	Glu	Ala	Val	Gly	Ile	Phe	Phe	Ala	Leu	Thr
	210					215					220				
Ile	Gly	Cys	Leu	Phe	Tyr	Leu	Gly	Met	Ile	Leu	Met	Leu	Asn	Pro	Leu
225					230				235						240
Tyr	Trp	Leu	Ser	Leu	Ala	Ile	Ala	Ile	Val	Gly	Trp	Val	Ile	Gln	Tyr
			245						250					255	
Ile	Gln	Leu	Ser	Ala	Pro	Thr	Pro	Glu	Pro	Lys	Leu	Tyr	Gly	Gln	Ile
			260					265					270		
Phe	Gly	Gln	Asn	Val	Ile	Ile	Gly	Phe	Val	Leu	Leu	Ala	Gly	Met	Leu
		275					280					285			
Leu	Gly	Trp	Leu												
			290												

&lt;210&gt; 33

&lt;211&gt; 316

&lt;212&gt; PRT

&lt;213&gt; Synechocystis sp

&lt;400&gt; 33

Met Val Thr Ser Thr Lys Ile His Arg Gln His Asp Ser Met Gly Ala  
 1 5 10 15  
 Val Cys Lys Ser Tyr Tyr Gln Leu Thr Lys Pro Arg Ile Ile Pro Leu  
 20 25 30  
 Leu Leu Ile Thr Thr Ala Ala Ser Met Trp Ile Ala Ser Glu Gly Arg  
 35 40 45  
 Val Asp Leu Pro Lys Leu Leu Ile Thr Leu Leu Gly Gly Thr Leu Ala  
 50 55 60  
 Ala Ala Ser Ala Gln Thr Leu Asn Cys Ile Tyr Asp Gln Asp Ile Asp  
 65 70 75 80  
 Tyr Glu Met Leu Arg Thr Arg Ala Arg Pro Ile Pro Ala Gly Lys Val  
 85 90 95  
 Gln Pro Arg His Ala Leu Ile Phe Ala Leu Ala Leu Gly Val Leu Ser  
 100 105 110  
 Phe Ala Leu Leu Ala Thr Phe Val Asn Val Leu Ser Gly Cys Leu Ala  
 115 120 125  
 Leu Ser Gly Ile Val Phe Tyr Met Leu Val Tyr Thr His Trp Leu Lys  
 130 135 140  
 Arg His Thr Ala Gln Asn Ile Val Ile Gly Gly Ala Ala Gly Ser Ile  
 145 150 155 160  
 Pro Pro Leu Val Gly Trp Ala Ala Val Thr Gly Asp Leu Ser Trp Thr  
 165 170 175  
 Pro Trp Val Leu Phe Ala Leu Ile Phe Leu Trp Thr Pro Pro His Phe  
 180 185 190  
 Trp Ala Leu Ala Leu Met Ile Lys Asp Asp Tyr Ala Gln Val Asn Val  
 195 200 205  
 Pro Met Leu Pro Val Ile Ala Gly Glu Glu Lys Thr Val Ser Gln Ile  
 210 215 220  
 Trp Tyr Tyr Ser Leu Leu Val Val Pro Phe Ser Leu Leu Leu Val Tyr  
 225 230 235 240  
 Pro Leu His Gln Leu Gly Ile Leu Tyr Leu Ala Ile Ala Ile Ile Leu  
 245 250 255  
 Gly Gly Gln Phe Leu Val Lys Ala Trp Gln Leu Lys Gln Ala Pro Gly  
 260 265 270  
 Asp Arg Asp Leu Ala Arg Gly Leu Phe Lys Phe Ser Ile Phe Tyr Leu  
 275 280 285  
 Met Leu Leu Cys Leu Ala Met Val Ile Asp Ser Leu Pro Val Thr His  
 290 295 300  
 Gln Leu Val Ala Gln Met Gly Thr Leu Leu Leu Gly  
 305 310 315

&lt;210&gt; 34

&lt;211&gt; 324

&lt;212&gt; PRT

&lt;213&gt; Synechocystis. sp

&lt;400&gt; 34

Met Ser Asp Thr Gln Asn Thr Gly Gln Asn Gln Ala Lys Ala Arg Gln  
 1 5 10 15  
 Leu Leu Gly Met Lys Gly Ala Ala Pro Gly Glu Ser Ser Ile Trp Lys  
 20 25 30  
 Ile Arg Leu Gln Leu Met Lys Pro Ile Thr Trp Ile Pro Leu Ile Trp  
 35 40 45  
 Gly Val Val Cys Gly Ala Ala Ser Ser Gly Gly Tyr Ile Trp Ser Val  
 50 55 60  
 Glu Asp Phe Leu Lys Ala Leu Thr Cys Met Leu Leu Ser Gly Pro Leu  
 65 70 75 80  
 Met Thr Gly Tyr Thr Gln Thr Leu Asn Asp Phe Tyr Asp Arg Asp Ile  
 85 90 95

Asp Ala Ile Asn Glu Pro Tyr Arg Pro Ile Pro Ser Gly Ala Ile Ser  
 100 105 110  
 Val Pro Gln Val Val Thr Gln Ile Leu Ile Leu Leu Val Ala Gly Ile  
 115 120 125  
 Gly Val Ala Tyr Gly Leu Asp Val Trp Ala Gln His Asp Phe Pro Ile  
 130 135 140  
 Met Met Val Leu Thr Leu Gly Gly Ala Phe Val Ala Tyr Ile Tyr Ser  
 145 150 155 160  
 Ala Pro Pro Leu Lys Leu Lys Gln Asn Gly Trp Leu Gly Asn Tyr Ala  
 165 170 175  
 Leu Gly Ala Ser Tyr Ile Ala Leu Pro Trp Trp Ala Gly His Ala Leu  
 180 185 190  
 Phe Gly Thr Leu Asn Pro Thr Ile Met Val Leu Thr Leu Ile Tyr Ser  
 195 200 205  
 Leu Ala Gly Leu Gly Ile Ala Val Val Asn Asp Phe Lys Ser Val Glu  
 210 215 220  
 Gly Asp Arg Gln Leu Gly Leu Lys Ser Leu Pro Val Met Phe Gly Ile  
 225 230 235 240  
 Gly Thr Ala Ala Trp Ile Cys Val Ile Met Ile Asp Val Phe Gln Ala  
 245 250 255  
 Gly Ile Ala Gly Tyr Leu Ile Tyr Val His Gln Gln Leu Tyr Ala Thr  
 260 265 270  
 Ile Val Leu Leu Leu Leu Ile Pro Gln Ile Thr Phe Gln Asp Met Tyr  
 275 280 285  
 Phe Leu Arg Asn Pro Leu Glu Asn Asp Val Lys Tyr Gln Ala Ser Ala  
 290 295 300  
 Gln Pro Phe Leu Val Phe Gly Met Leu Ala Thr Gly Leu Ala Leu Gly  
 305 310 315 320  
 His Ala Gly Ile

&lt;210&gt; 35

&lt;211&gt; 307

&lt;212&gt; PRT

&lt;213&gt; Synechocystis sp

&lt;400&gt; 35

Met Thr Glu Ser Ser Pro Leu Ala Pro Ser Thr Ala Pro Ala Thr Arg  
 1 5 10 15  
 Lys Leu Trp Leu Ala Ala Ile Lys Pro Pro Met Tyr Thr Val Ala Val  
 20 25 30  
 Val Pro Ile Thr Val Gly Ser Ala Val Ala Tyr Gly Leu Thr Gly Gln  
 35 40 45  
 Trp His Gly Asp Val Phe Thr Ile Phe Leu Leu Ser Ala Ile Ala Ile  
 50 55 60  
 Ile Ala Trp Ile Asn Leu Ser Asn Asp Val Phe Asp Ser Asp Thr Gly  
 65 70 75 80  
 Ile Asp Val Arg Lys Ala His Ser Val Val Asn Leu Thr Gly Asn Arg  
 85 90 95  
 Asn Leu Val Phe Leu Ile Ser Asn Phe Phe Leu Leu Ala Gly Val Leu  
 100 105 110  
 Gly Leu Met Ser Met Ser Trp Arg Ala Gln Asp Trp Thr Val Leu Glu  
 115 120 125  
 Leu Ile Gly Val Ala Ile Phe Leu Gly Tyr Thr Tyr Gln Gly Pro Pro  
 130 135 140  
 Phe Arg Leu Gly Tyr Leu Gly Leu Gly Glu Leu Ile Cys Leu Ile Thr  
 145 150 155 160  
 Phe Gly Pro Leu Ala Ile Ala Ala Ala Tyr Tyr Ser Gln Ser Gln Ser  
 165 170 175  
 Phe Ser Trp Asn Leu Leu Thr Pro Ser Val Phe Val Gly Ile Ser Thr  
 180 185 190

Ala Ile Ile Leu Phe Cys Ser His Phe His Gln Val Glu Asp Asp Leu  
 195 200 205  
 Ala Ala Gly Lys Lys Ser Pro Ile Val Arg Leu Gly Thr Lys Leu Gly  
 210 215 220  
 Ser Gln Val Leu Thr Leu Ser Val Val Ser Leu Tyr Leu Ile Thr Ala  
 225 230 235 240  
 Ile Gly Val Leu Cys His Gln Ala Pro Trp Gln Thr Leu Leu Ile Ile  
 245 250 255  
 Ala Ser Leu Pro Trp Ala Val Gln Leu Ile Arg His Val Gly Gln Tyr  
 260 265 270  
 His Asp Gln Pro Glu Gln Val Ser Asn Cys Lys Phe Ile Ala Val Asn  
 275 280 285  
 Leu His Phe Phe Ser Gly Met Leu Met Ala Ala Gly Tyr Gly Trp Ala  
 290 295 300  
 Gly Leu Gly  
 305

<210> 36  
 <211> 927  
 <212> DNA  
 <213> Synechocystis sp

<400> 36  
 atggcaacta tccaagcttt ttggcgcttc tcccgccccc ataccatcat tgggtacaact  
 60  
 ctgagcgtct gggctgtgta tctgttaact attctcgggg atggaaactc agttaactcc  
 120  
 cctgcttccc tggatttagt gttcggcgct tggctggcct gcctgttggg taatgtgtac  
 180  
 attgtcggcc tcaaccaatt gtgggatgtg gacattgacc gcatcaataa gccgaatttg  
 240  
 cccctagcta acggagattt ttctatcgcc cagggccggt ggattgtggg actttgtggc  
 300  
 gttgcttcct tggcgatcgc ctggggatta gggctatggc tggggctaac ggtgggcatt  
 360  
 agtttgatta ttggcacggc ctattcgggt cgcagtgga ggttaaagcg cttttccctg  
 420  
 ctggcgcccc tgtgtattct gacgggtgcg ggaattgtgg ttaacttggg cttattttta  
 480  
 ttttttagaa ttggtttagg ttatcccccc actttaataa ccccatctg ggttttgact  
 540  
 ttatttatct tagttttcac cgtggcgatc gccattttta aagatgtgcc agatatggaa  
 600  
 ggcgatcggc aatttaagat tcaaacttta actttgcaaa tcggcaaaca aaacgttttt  
 660  
 cggggaacct taattttact cactggttgt tatttagcca tggcaatctg gggcttatgg  
 720  
 gcggctatgc ctttaaatac tgctttcttg attgtttccc atttgtgctt attagcctta  
 780  
 ctctgggtggc ggagtcgaga tgtacactta gaaagcaaaa ccgaaattgc tagtttttat  
 840  
 cagtttatct ggaagctatt tttcttagag tacttgctgt atcccttggc tctgtgggta  
 900  
 cctaattttt ctaatactat ttttttag  
 927

<210> 37  
 <211> 308  
 <212> PRT  
 <213> Synechocystis sp

&lt;400&gt; 37

Met Ala Thr Ile Gln Ala Phe Trp Arg Phe Ser Arg Pro His Thr Ile  
 1 5 10 15  
 Ile Gly Thr Thr Leu Ser Val Trp Ala Val Tyr Leu Leu Thr Ile Leu  
 20 25 30  
 Gly Asp Gly Asn Ser Val Asn Ser Pro Ala Ser Leu Asp Leu Val Phe  
 35 40 45  
 Gly Ala Trp Leu Ala Cys Leu Leu Gly Asn Val Tyr Ile Val Gly Leu  
 50 55 60  
 Asn Gln Leu Trp Asp Val Asp Ile Asp Arg Ile Asn Lys Pro Asn Leu  
 65 70 75 80  
 Pro Leu Ala Asn Gly Asp Phe Ser Ile Ala Gln Gly Arg Trp Ile Val  
 85 90 95  
 Gly Leu Cys Gly Val Ala Ser Leu Ala Ile Ala Trp Gly Leu Gly Leu  
 100 105 110  
 Trp Leu Gly Leu Thr Val Gly Ile Ser Leu Ile Ile Gly Thr Ala Tyr  
 115 120 125  
 Ser Val Pro Pro Val Arg Leu Lys Arg Phe Ser Leu Leu Ala Ala Leu  
 130 135 140  
 Cys Ile Leu Thr Val Arg Gly Ile Val Val Asn Leu Gly Leu Phe Leu  
 145 150 155 160  
 Phe Phe Arg Ile Gly Leu Gly Tyr Pro Pro Thr Leu Ile Thr Pro Ile  
 165 170 175  
 Trp Val Leu Thr Leu Phe Ile Leu Val Phe Thr Val Ala Ile Ala Ile  
 180 185 190  
 Phe Lys Asp Val Pro Asp Met Glu Gly Asp Arg Gln Phe Lys Ile Gln  
 195 200 205  
 Thr Leu Thr Leu Gln Ile Gly Lys Gln Asn Val Phe Arg Gly Thr Leu  
 210 215 220  
 Ile Leu Leu Thr Gly Cys Tyr Leu Ala Met Ala Ile Trp Gly Leu Trp  
 225 230 235 240  
 Ala Ala Met Pro Leu Asn Thr Ala Phe Leu Ile Val Ser His Leu Cys  
 245 250 255  
 Leu Leu Ala Leu Leu Trp Trp Arg Ser Arg Asp Val His Leu Glu Ser  
 260 265 270  
 Lys Thr Glu Ile Ala Ser Phe Tyr Gln Phe Ile Trp Lys Leu Phe Phe  
 275 280 285  
 Leu Glu Tyr Leu Leu Tyr Pro Leu Ala Leu Trp Leu Pro Asn Phe Ser  
 290 295 300  
 Asn Thr Ile Phe  
 305

&lt;210&gt; 38

&lt;211&gt; 1092

&lt;212&gt; DNA

&lt;213&gt; Synechocystis sp

&lt;400&gt; 38

atgaaatttc cgccccacag tggttacat tggcaaggct aatcaccttt ctttgaagg  
 60  
 tggtagctgc gcctgctttt gccccaatcc ggggaaagtt ttgcttttat gtactccatc  
 120  
 gaaaatcctg ctagcgatca tcattacggc ggcgggtgctg tgcaaatttt agggccggct  
 180  
 acgaaaaaac aagaaaatca ggaagaccaa cttgtttggc ggacatttcc ctcggtaaaa  
 240  
 aaattttggg ccagtcctcg ccagtttgcc ctagggcatt ggggaaaatg tagggataac  
 300  
 aggcaggcga aaccctact ctccgaagaa ttttttgcca cggtcaagga aggttatcaa  
 360  
 atccatcaaa atcagcacca aggacaaatc attcatggcg atcgccattg tcgttggcag



420  
 ttcaccgtag aaccggaagt aacttggggg agtcctaacc gatttcctcg ggctacagcg  
 480  
 ggttggcttt cctttttacc cttgtttgat cccggttggc aaattctttt agcccaaggt  
 540  
 agagcgacg gctggctgaa atggcagagg gaacagtatg aatttgacca cgccctagtt  
 600  
 tatgccgaaa aaaattgggg tcactccttt ccctcccgtt ggttttggct ccaagcaaat  
 660  
 tattttcctg accatccagg actgagcgtc actgccgctg gcggggaacg gattgttctt  
 720  
 ggtcgccccg aagaggtagc ttttaattggc ttacatcacc aaggtaattt ttacgaattt  
 780  
 ggcccggggc atggcacagt cacttggcaa gtagctccct ggggccggtt gcaattaaaa  
 840  
 gccagcaatg ataggtattg ggtcaagttg tccggaaaaa cagataaaaa aggcagttta  
 900  
 gtccacactc ccaccgcca gggcttacia ctcaactgcc gagataccac taggggctat  
 960  
 ttgtatttgc aattgggatc tgtgggtcac ggcctgatag tgcaagggga aacggacacc  
 1020  
 gcggggctag aagttggagg tgattggggt ttaacagagg aaaatttgag caaaaaaaca  
 1080  
 gtgccattct ga  
 1092

<210> 39  
 <211> 363  
 <212> PRT  
 <213> Synechocystis sp

<400> 39  
 Met Lys Phe Pro Pro His Ser Gly Tyr His Trp Gln Gly Gln Ser Pro  
 1 5 10 15  
 Phe Phe Glu Gly Trp Tyr Val Arg Leu Leu Leu Pro Gln Ser Gly Glu  
 20 25 30  
 Ser Phe Ala Phe Met Tyr Ser Ile Glu Asn Pro Ala Ser Asp His His  
 35 40 45  
 Tyr Gly Gly Gly Ala Val Gln Ile Leu Gly Pro Ala Thr Lys Lys Gln  
 50 55 60  
 Glu Asn Gln Glu Asp Gln Leu Val Trp Arg Thr Phe Pro Ser Val Lys  
 65 70 75 80  
 Lys Phe Trp Ala Ser Pro Arg Gln Phe Ala Leu Gly His Trp Gly Lys  
 85 90 95  
 Cys Arg Asp Asn Arg Gln Ala Lys Pro Leu Leu Ser Glu Glu Phe Phe  
 100 105 110  
 Ala Thr Val Lys Glu Gly Tyr Gln Ile His Gln Asn Gln His Gln Gly  
 115 120 125  
 Gln Ile Ile His Gly Asp Arg His Cys Arg Trp Gln Phe Thr Val Glu  
 130 135 140  
 Pro Glu Val Thr Trp Gly Ser Pro Asn Arg Phe Pro Arg Ala Thr Ala  
 145 150 155 160  
 Gly Trp Leu Ser Phe Leu Pro Leu Phe Asp Pro Gly Trp Gln Ile Leu  
 165 170 175  
 Leu Ala Gln Gly Arg Ala His Gly Trp Leu Lys Trp Gln Arg Glu Gln  
 180 185 190  
 Tyr Glu Phe Asp His Ala Leu Val Tyr Ala Glu Lys Asn Trp Gly His  
 195 200 205  
 Ser Phe Pro Ser Arg Trp Phe Trp Leu Gln Ala Asn Tyr Phe Pro Asp  
 210 215 220  
 His Pro Gly Leu Ser Val Thr Ala Ala Gly Gly Glu Arg Ile Val Leu

225					230					235					240
Gly	Arg	Pro	Glu	Glu	Val	Ala	Leu	Ile	Gly	Leu	His	His	Gln	Gly	Asn
				245					250					255	
Phe	Tyr	Glu	Phe	Gly	Pro	Gly	His	Gly	Thr	Val	Thr	Trp	Gln	Val	Ala
			260					265					270		
Pro	Trp	Gly	Arg	Trp	Gln	Leu	Lys	Ala	Ser	Asn	Asp	Arg	Tyr	Trp	Val
		275					280				285				
Lys	Leu	Ser	Gly	Lys	Thr	Asp	Lys	Lys	Gly	Ser	Leu	Val	His	Thr	Pro
	290					295					300				
Thr	Ala	Gln	Gly	Leu	Gln	Leu	Asn	Cys	Arg	Asp	Thr	Thr	Arg	Gly	Tyr
305					310					315					320
Leu	Tyr	Leu	Gln	Leu	Gly	Ser	Val	Gly	His	Gly	Leu	Ile	Val	Gln	Gly
			325					330						335	
Glu	Thr	Asp	Thr	Ala	Gly	Leu	Glu	Val	Gly	Gly	Asp	Trp	Gly	Leu	Thr
			340					345					350		
Glu	Glu	Asn	Leu	Ser	Lys	Lys	Thr	Val	Pro	Phe					
		355					360								

&lt;210&gt; 40

&lt;211&gt; 56

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: Oligonucleotide

&lt;400&gt; 40

cgcgatttaa atggcgcgcc ctgcaggcgg ccgcctgcag ggcgcgccat ttaaat  
56

&lt;210&gt; 41

&lt;211&gt; 32

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: Oligonucleotide

&lt;400&gt; 41

tgcgaggatcc gcggccgcaa gcttcctgca gg  
32

&lt;210&gt; 42

&lt;211&gt; 32

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: Oligonucleotide

&lt;400&gt; 42

tcgacctgca ggaagcttgc ggccgcggat cc  
32

&lt;210&gt; 43

&lt;211&gt; 32

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: Oligonucleotide

<400> 43

tcgacctgca ggaagcttgc ggccgcggat cc  
32

<210> 44

<211> 32

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Oligonucleotide

<400> 44

tcgaggatcc gcggccgcaa gcttcctgca gg  
32

<210> 45

<211> 36

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Oligonucleotide

<400> 45

tcgaggatcc gcggccgcaa gcttcctgca ggagct  
36

<210> 46

<211> 28

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Oligonucleotide

<400> 46

cctgcaggaa gcttgcggcc gcggatcc  
28

<210> 47

<211> 36

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Oligonucleotide

<400> 47

tcgacctgca ggaagcttgc ggccgcggat ccagct  
36

<210> 48

<211> 28

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Oligonucleotide

<400> 48

ggatccgcgg ccgcaagctt cctgcagg  
28

<210> 49

<211> 39

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Oligonucleotide

<400> 49

gatcacctgc aggaagcttg cggccgcgga tccaatgca  
39

<210> 50

<211> 31

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Oligonucleotide

<400> 50

ttggatccgc ggccgcaagc ttcctgcagg t  
31

<210> 51

<211> 41

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Oligonucleotide

<400> 51

ggatccgcgg ccgcacaatg gagtctctgc tctctagttc t  
41

<210> 52

<211> 38

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Oligonucleotide

<400> 52

ggatcctgca ggtcacttca aaaaaggtaa cagcaagt  
38

<210> 53

<211> 45

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Oligonucleotide

<400> 53

ggatccgcgg ccgcacaatg gcgttttttg ggctctcccg tgttt  
45

<210> 54  
<211> 40  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Oligonucleotide

<400> 54  
ggatcctgca ggattattgaa aacttcttcc aagtacaact  
40

<210> 55  
<211> 38  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Oligonucleotide

<400> 55  
ggatccgcgg ccgcacaatg tggcgaagat ctgttggt  
38

<210> 56  
<211> 37  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Oligonucleotide

<400> 56  
ggatcctgca ggatcatggag agtagaagga aggagct  
37

<210> 57  
<211> 50  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Oligonucleotide

<400> 57  
ggatccgcgg ccgcacaatg gtacttgccg aggttccaaa gcttgctct  
50

<210> 58  
<211> 38  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Oligonucleotide

<400> 58  
ggatcctgca ggtcacttgt ttctggatgat gactctat

38

<210> 59  
<211> 38  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Oligonucleotide

<400> 59  
ggatccgcgg ccgcacaatg acttcgattc tcaacact  
38

<210> 60  
<211> 36  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Oligonucleotide

<400> 60  
ggatcctgca ggtcagtgtt gcgatgctaa tgccgt  
36

<210> 61  
<211> 22  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Oligonucleotide

<400> 61  
taatgtgtac attgtcggcc tc  
22

<210> 62  
<211> 60  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Oligonucleotide

<400> 62  
gcaatgtaac atcagagatt ttgagacaca acgtggcttt ccacaattcc ccgcaccgtc  
60

<210> 63  
<211> 22  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Oligonucleotide

<400> 63  
aggctaataa gcacaaatgg ga  
22



<210> 64  
<211> 63  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Oligonucleotide

<400> 64  
ggtatgagtc agcaacacct tcttcacgag gcagacctca gcggaattgg tttaggttat  
60  
ccc  
63

<210> 65  
<211> 26  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Oligonucleotide

<400> 65  
ggatccatgg ttgcccaaac cccatc  
26

<210> 66  
<211> 61  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Oligonucleotide

<400> 66  
gcaatgtaac atcagagatt ttgagacaca acgtggcctt gggtaagcaa caatgaccgg  
60  
c  
1

6

<210> 67  
<211> 25  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Oligonucleotide

<400> 67  
gaattctcaa agccagccca gtaac  
25

<210> 68  
<211> 63  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Oligonucleotide

<400> 68  
ggatgagtc agcaacacct tcttcacgag gcagacctca gcgggtgcga aaagggtttt  
60  
ccc  
63

<210> 69  
<211> 23  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Oligonucleotide

<400> 69  
ccagtggttt aggctgtgtg gtc  
23

<210> 70  
<211> 21  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Oligonucleotide

<400> 70  
ctgagttgga tgtattggat c  
21

<210> 71  
<211> 28  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Oligonucleotide

<400> 71  
ggatccatgg ttacttcgac aaaaatcc  
28

<210> 72  
<211> 60  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Oligonucleotide

<400> 72  
gcaatgtaac atcagagatt ttgagacaca acgtggcttt gctaggcaac cgcttagtac  
60

<210> 73  
<211> 28  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Oligonucleotide

<400> 73  
gaattcttaa cccaacagta aagttccc  
28

<210> 74  
<211> 63  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Oligonucleotide

<400> 74  
ggtatgagtc agcaacacct tcttcacgag gcagacctca gcgccggcat tgtcttttac  
60  
atg  
63

<210> 75  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Oligonucleotide

<400> 75  
ggaacccttg cagccgcttc  
20

<210> 76  
<211> 22  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Oligonucleotide

<400> 76  
gtatgcccaa ctggtgcaga gg  
22

<210> 77  
<211> 28  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Oligonucleotide

<400> 77  
ggatccatgt ctgacacaca aaataccg  
28

<210> 78  
<211> 62  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Oligonucleotide

<400> 78

gcaatgtaac atcagagatt ttgagacaca acgtggcttt cgccaatacc agccaccaac  
60

ag

62

<210> 79

<211> 27

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Oligonucleotide

<400> 79

gaattctcaa atccccgcat ggcctag  
27

<210> 80

<211> 65

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Oligonucleotide

<400> 80

ggtatgagtc agcaacacct tcttcacgag gcagacctca gcggcctacg gcttggacgt  
60

gtggg

65

<210> 81

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Oligonucleotide

<400> 81

cacttggatt cccctgatct g  
21

<210> 82

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Oligonucleotide

<400> 82

gcaatacccg cttggaaaac g  
21

<210> 83

<211> 29

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Oligonucleotide

<400> 83

ggatccatga ccgaatcttc gcccttagc

29

<210> 84

<211> 61

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Oligonucleotide

<400> 84

gcaatgtaac atcagagatt ttgagacaca acgtggcttt caatcctagg tagccgaggc

60

g

1

6

<210> 85

<211> 27

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Oligonucleotide

<400> 85

gaattcttag cccaggccag cccagcc

27

<210> 86

<211> 66

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Oligonucleotide

<400> 86

ggtatgagtc agcaacacct tcttcacgag gcagacctca gcggggaatt gatttggtta

60

attacc

66

<210> 87

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Oligonucleotide

<400> 87

gcgatcgcca ttatcgcttg g

21

<210> 88  
<211> 24  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Oligonucleotide

<400> 88  
gcagactggc aattatcagt aacg  
24

<210> 89  
<211> 25  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Oligonucleotide

<400> 89  
ccatggattc gagtaaagtt gtcgc  
25

<210> 90  
<211> 25  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Oligonucleotide

<400> 90  
gaattcactt caaaaaaggt aacag  
25

<210> 91  
<211> 4550  
<212> DNA  
<213> Arabidopsis sp

<400> 91  
attttacacc aatttgatca cttaactaaa ttaattaaat tagatgatta tcccaccata  
60  
tttttgagca ttaaaccata aaaccatagt tataagtaac tgttttaac gaatatgact  
120  
cgattaagat taggaaaaat ttataaccgg taattaagaa aacattaacc gtagtaaccg  
180  
taaatgccga ttcctccctt gtctaaaaga cagaaaacat atattttatt ttgccccata  
240  
tgtttcactc tatttaattt caggcacaat acttttggtt ggtaacaaaa ctaaaagga  
300  
caacacgtga tacttttcct cgtccgtcag tcagattttt tttaaactag aaacaagtgg  
360  
caaactaca ccacattttt tgcttaatct attaacttgt aagtttttaa ttcctaaaaa  
420  
agtctaacta attcttctaa tataagtaca ttccctaaat ttcccaaaaa gtcaaattaa  
480  
taattttcaa aatctaattt aaatatctaa taattcaaaa tcattaaaaa gacacgcaac  
540  
aatgacacca attaatcatc ctcgaccac acaattctac agttctcatg cttaaccata



600  
ttttttgtct tctgttcctt caaaatcatt tctttctctt ctttgattcc caaagatcac  
660  
ttctttgtct ttgatttttg attttttttc tctctggcgt gaaggaagaa gctttatttc  
720  
atggagtctc tgctctctag ttcttctctt gtttccgctg gtaaactctg tccttttctg  
780  
gtttcagggt ttatttggtg tttagggttc gtttttgbga ttcagaacca tacaaaaagt  
840  
ttgaactttt ctgaatataa aataaggaaa aagtttcgat ttttataatg aattgtttac  
900  
tagatcgaag taggtgacaa aggttattgt gtggagaagc ataatttctg ggcttgactt  
960  
tgaattttgt ttctcatgca tgcaacttat caatcagctg gtgggttttg ttggaagaag  
1020  
cagaatctaa agctccactc tttatcagggt tcgttaggggt tttatgggtt tttgaaatta  
1080  
aatactcaat catcttagtc tcattattct attggttgaa tcacattttc taatttgga  
1140  
tttatgagac aatgtatgtt ggacttaggt gaagttcttc tctttgggta tagttgaagt  
1200  
gttactgatg ttgttttagct ctttacacca atatatacac ccaattttgc agaaatccga  
1260  
gttctgcgtt gtgattcgag taaagttgtc gcaaaaccga agtttaggaa caatcttggt  
1320  
aggcctgatg gtcaaggatc ttcattgttg ttgtatccaa aacataagtc gagatttcgg  
1380  
gttaatgcca ctgcgggtca gcctgaggct ttcgactcga atagcaaaca gaagtctttt  
1440  
agagactcgt tagatgcgtt ttacagggtt tctaggectc atacagttat tggcacagtt  
1500  
aagtttctct ttaaaaatgt aactctttta aaacgcaatc tttcagggtt ttcaaggaga  
1560  
taacattagc tctgtgattg gatttgcagg tgcttagcat tttatctgta tctttcttag  
1620  
cagtagagaa ggtttctgat atatctcctt tacttttcac tggcatcttg gaggtaatga  
1680  
atatataaca cataatgacc gatgaagaag atacattttt ttcgtctctc tgttttaaca  
1740  
attgggtttt gttttcaggc tgttgttgca gctctcatga tgaacattta catagttggg  
1800  
ctaaatcagt tgtctgatgt tgaaatagat aaggtaacat gcaaattttc ttcatatgag  
1860  
ttcgagagac tgatgagatt aatagcagct agtgcctaga tcatctctat gtgggttttt  
1920  
gcaggttaac aagccctatc ttccattggc atcaggagaa tattctgtta acaccggcat  
1980  
tgcaatagta gcttccttct ccatcatggg atgggtgccat tttcacaaaa tttcaacttt  
2040  
tagaattcta taagttactg aaatagtttg ttataaatcg ttatagagtt tctggcttgg  
2100  
gtggattggt ggttcatggc cattgttctg ggctcttttt gtgagtttca tgctcggtac  
2160  
tgcatactct atcaatgtaa gtaagtttct caatactaga atttggctca aatcaaaatc  
2220  
tgcagtttct agtttttaggt taatgagggt ttaataactt acttctacta caaacagttg  
2280  
ccacttttac ggtggaaaag atttgcattg gttgcagcaa tgtgtatcct cgctgtccga  
2340  
gctattattg ttcaaatcgc cttttatcta catattcagg tactaaacca ttttccttat  
2400

gtttttagt tgttttcatc aaaatcactt ttatattact aaagctgtga aactttgttg  
2460  
cagacacatg tgtttggaag accaatcttg ttcactaggc ctcttatttt cgccactgcg  
2520  
tttatgagct ttttctctgt cgttattgca ttgtttaagg taaacaaaga tggaaaaaga  
2580  
ttaaatctat gtatacttaa agtaaagcat tctactgtta ttgatgagaa gttttctttt  
2640  
ttggttggaat gcaggatata cctgatatcg aaggggataa gatattcgga atccgatcat  
2700  
tctctgtaac tctgggtcag aaacgggtac gatattctaaa ctaaagaaat tgttttgact  
2760  
caagtgttgg attaagatta cagaagaaag aaaactgttt ttgtttcttg caaaattcag  
2820  
gtgttttggga catgtgttac actacttcaa atggcttacg ctgttgcaat tctagttgga  
2880  
gccacatctc cattcatatg gagcaaagtc atctcggtaa caatctttct ttacccatcg  
2940  
aaaactcgct aattcatcgt ttgagtggta ctggtttcat tttgttcogt tctgttgatt  
3000  
ttttttcagg ttgtgggtca tgttatactc gcaacaactt tgtgggctcg agctaagtcc  
3060  
gttgatctga gtagcaaaac cgaaataact tcatgttata tgttcatatg gaaggttaga  
3120  
ttcgtttata aatagagtct ttactgcctt tttatgcgct ccaatttgga attaaaatag  
3180  
cctttcagtt tcatcgaatc accattatac tgataaattc tcatttctgc atcagctctt  
3240  
ttatgcagag tacttgctgt tacctttttt gaagtgactg acattagaag agaagaagat  
3300  
ggagataaaa gaataagtca tcactatgct tctgttttta ttacaagttc atgaaattag  
3360  
gtagtgaact agtgaattag agttttatct tgaacatgg cagactgcaa aaatatgtca  
3420  
aagatatgaa tttctgttgg gtaaagaagt ctctgcttgg gcaaaatctt aaggttcggg  
3480  
gtgttgatat aatgctaagc gaagaaatcg attctatgta gaaatttccg aaactatgtg  
3540  
taaacatgtc agaacatctc cattctatat cttcttctgc aagaaagctc tgtttttate  
3600  
acctaaactc tttatctctg tgtagttaag atatgtatat gtacgtgact acattttttt  
3660  
gttgatgtaa tttgcagaac gtatggattt ttgttagaaa gcatgagttc gaaagtatat  
3720  
gtttatatat atggataatt cagacctaac gtcgaagctc acaagcataa attcactact  
3780  
atagtttgct ctgtaataga tagttccatt gatgtcttga aactgtacgt aactgcctgg  
3840  
gcgttttgtg gttgatactg actactgagt gttctttgtg agtgttgtaa gtatacaaga  
3900  
agaagaatat aggctcacgg gaacgactgt ggtggaagat gaaatggaga tcatcacgta  
3960  
gcggctttgc caaagaccga gtcacgatcg agtctatgaa gtctttacag ctgctgatta  
4020  
tgattgacca ttgcttagag acgcattgga atcttactag ggacttgcct gggagtttct  
4080  
tcaagtacgt gtcagatcat acgatgtagg agatttcacg gctttgatgt gtttggttgg  
4140  
agtcacaatg cttaatgggc ttattggccc aataatagct agctcttttg ctttagccgt  
4200  
ttcgtttgtc ccctgggtgg gagtattatt agggatgggt gtgaccaaag tcaccagacc

4260  
tagagtgaat ctagtagagt cctagaccat ggtccatggc ttttatttgt aatttgaaaa  
4320  
atgaacaatt ctttttgtaa ggaaaacttt tatatagtag acgtttacta tatagaaact  
4380  
agttgaacta acttcgtgca attgcataat aatgggtgtga aatagagggt gcaaaactca  
4440  
ataaacattt cgacgtacca agagttcgaa acaataagca aaatagattt ttttgcttca  
4500  
gactaatttg tacaatgaat ggttaataaa ccattgaagc ttttattaat  
4550

<210> 92  
<211> 4450  
<212> DNA  
<213> Arabidopsis sp

<400> 92  
tttaggttac aaaatcaatg atattgcgta tgtcaactat aaaagccaaa agtaaagcct  
60  
cttgtttgac cagaagggtca tgatcattgt atacatacag ccaaactacc tcctggaaga  
120  
aaagacatgg atcccaaaca acaacaatag cttcttttac aagaaccagt agtaactagt  
180  
cactaatcta aaagagttaa gtttcagctt ttctggcaat ggctccttga tcatttcaat  
240  
cctgaaggag acccactttg tagcaagacc atgtcctctg tttcacttac agtgtgtctc  
300  
aaaagtctac ttcaattctt catatatagg ttcttcacac tacagcttca tcctcattcg  
360  
ttgacagaga gagagtcttt attgaaaact tcttccaagt acaactccac taaatataat  
420  
agcaccaaac cacttgttcg acacaaatct gtacagatat aaaaacacta ttaggttttc  
480  
caaggcaaat cacataattg gattgtgaaa gagtacaaaa gataaaccce aattttcata  
540  
ctttctactg cagtcagcac cagatgataa gtcagctgtc cctatttgcc atcctaactg  
600  
tcctgatgca ggggccagtg atgcgtaata ttgccaccct taatcattag agcgagaaac  
660  
aaaaagaatc aaaagacagt aaatggaatt aggaatcaca aatgagtcct tgtaaagttt  
720  
attgagtacc gagatctgca ctgaatccag aaagtgcag aaaacctatg gatgctgtgc  
780  
caaatccagt taaccaaagc tttgtattat caccgaatct aagggtgtgt gacttaacac  
840  
caacttttac atcatcttct ttgtcctgga gacacaatat attagacatt agtccatgga  
900  
aaaaaaatga tttaacctag aatatctcaa aattacttgc ataaaaactg aacttgagct  
960  
gaaatttttg gttcgtagct tgtggcatat actatttcat tttcaatggg ccacaaaggt  
1020  
aactttcttt tctcacttct gttgcaaagc ggaagacttt tatggggcta actcttcact  
1080  
taaagtatag aaatcagatg gaaaagggtg gagatcaggg taattttctt ctttatgatt  
1140  
gacaaaagtc gaacatcgaa atggatgcat ttgcatgaga catgaaacaa aagctgaaaa  
1200  
agaaatctgt ggtggtgaag ctagaaaaag aaaacaaagc aagcaatatg cacacattga  
1260  
gattaactac tttgctactg gtcataatca aatagatttt gaagctaaaa aataaaaagt

1320  
gaatatacct gatgtgcata aatagtatca taaacaagg tccagcagac tccggagaga  
1380  
tagagagggga gtacaataga tgggtgctatg cttccttttaa ctgcagtcca tcctaacaat  
1440  
gctccccagt ttatgggtcaa acctaaaaag gcttgaggct gcaattataa aaacgaatca  
1500  
atcataagaa aatcagaaaa tatataatgt ctaactttga gaagccagaa tagattttaa  
1560  
ttacccaaaa tgtaaacctc ttcataagtg ggtaggaaaa gacaagtaac aaagatgaag  
1620  
cccctaaaac acggctgcag aatatacata ctgaaatgag ctcaagtaga aaagaatttg  
1680  
atcacaaaac taaagacaag acctgagaac atatcttcag aatttgggcc aactacataa  
1740  
gggtgaacca tatgtgtatg tgaattttta aacaaacact tgcaaatacg cgactttagg  
1800  
gcaagtaaaa aatccaaaca aacctgtaat tgtaagtgtg gagaagaatc cctaagccta  
1860  
aaagcaactg cagcccgaga aatccaatcc cttgaaatgg tgtcaaaaga ccactggcga  
1920  
taggtcttag ttttgtacga tcaacctgga tataaaagaa atttgtaaga caacataatc  
1980  
taaaacaaaa caaccatata aaatcttgag ctttacatac aagcaacca tctttgttta  
2040  
tggaagaatg aatccagtta catgaatgct gtgtatctac cctaactact aaacacatat  
2100  
ttcaatcgaa aaacatatcc caccttcacc atatctaaca cctgaagtct ttcacttttt  
2160  
gaacgaagtc atcagaacat gcagataagc tattacccaa aacagagata tgactggaaa  
2220  
tggtgtcgta aattgatcca acatagaaaa atcaagacca gttccagatg tcaaagcaat  
2280  
aacactttcc caccatgggt acagaaacca tagttacaca aaacatgttt cctaaaccaa  
2340  
catactaaag ggatatataa atttgacatc actttatcac cataccataa gatagcttaa  
2400  
aaacaaactg acctttgtat ctatgtcctg atcaagcaga tcatttatag tacaaccagc  
2460  
acctctaaga agtaatgctc cgcaacccaa taaagccata tatttaaaac ttggaaggct  
2520  
tccaggatca gcagccaacg caatcgacct atacaacaat gatggagatt cagagtatcg  
2580  
atctatttac atagctctgg aactagatcc atgacgaaac atggaacatc gttataatat  
2640  
ctaaagactt ccaaacagat tcctgagtaa gaaaccagc ggaactatag tactgtaaca  
2700  
tatataaaat caaagaaaac tcagggttat agcattatcc aatcctgatt tctgccaatc  
2760  
cttaaccact ctcccatgct atcaaaaacc tcagctcaag atcatactac ctaattgcct  
2820  
atgagctctt gggaagatca ttatggattt gataactgaa aaaagtaaca gagaaatagc  
2880  
agactgcaag aactactcca aacttctcca ctgatatgta tgtagtctaa caataataaa  
2940  
cagacataaa ttctttttatc aagcttcaag agcaagttag tcagaaaaca tcacagccaa  
3000  
accaaccagg aaaacacata actttatcac ataaaactaa atttaatgta atctgactta  
3060  
acataaacca tcctttggga cgaaaggaaa ctatataaac atgcagtctt tctttccctc  
3120

agctattctt tcggatggat tataatgaat ctcaaaagtg aaatgtcttg attctcagct  
 3180  
 acattactca aaggcgaaga taaacttacc acatacaagg ccacgcaagc aaccaagttc  
 3240  
 caatggggtt atccaatcga gcaagcttag cataacctct aacttcttct ggtaaataca  
 3300  
 aatctatcca agaagcttcc ttaacaacaa caccatcact cttctcctta tcatctttct  
 3360  
 tcggctttcc ctccaaaacc gaagaagacg acgacattcc acaaattaat ctgtaattcc  
 3420  
 aaccaacacc aaaaaacttc tcctgatgca attctcttcc ttacttccat acttggtaat  
 3480  
 tatcattcca tgaaggataa cacttagtga aaggatttgt gtaatgggta gtcacaggat  
 3540  
 tggacaagga tttatgttgt gattgcaaaa gagcagagga agaagatgga gttacggaga  
 3600  
 cggaagattt caacaaccgt cttgaaacac gggagagccc aaaaaacgcc atctttgaga  
 3660  
 gaaattgttg cctggaagaa acaaagactt gagatttcaa acgtaagtga attcttacga  
 3720  
 acgaaagcta acttctcaag agaatcagat tagtgattcc tcaaaaacaa aaaaaactat  
 3780  
 ctaatttcag tttcgagtga tgaagcctta agaatctaga acctccatgg cgtttcta  
 3840  
 ctctcagaga taatcgaatt ccttaaacia tcaaagctta gaaagagaag aacaacaaca  
 3900  
 acaacaaaaa aaatcagatt aacaaccgac cagagagcaa cgacgacgcc ggcgagaaag  
 3960  
 agcacgtcgt ctcgagcaa gacttcttct ccagtaacce ggatggatcg ttaatgggcc  
 4020  
 tgtagattat tatatttggg ccgaaacaat tgggtcagca aaaacttggg ggataatgaa  
 4080  
 gaaacacgta cagtatgcat ttaggtcca aattaattgg ccatataatt cgaatcagat  
 4140  
 aaactaatca accctacct tacttatttc tcaactgttt tatttctacc ttagtagttg  
 4200  
 aagaaacact tttatttatc ttttcgggac ccaaatttga taggatcggg ccattactca  
 4260  
 tgagcgtcag acacatatta gccttatcag attagtgggg taagggtttt ttaattcggg  
 4320  
 aagaagcaac aatcaatgtc ggagaaatta aagaatctgc atgggcgtgg cgtgatgata  
 4380  
 tgtgcatatg gagtcagttg ccgatcatat ataactattt ataaactaca tataaagact  
 4440  
 actaatagat  
 4450

&lt;210&gt; 93

&lt;211&gt; 2850

&lt;212&gt; DNA

&lt;213&gt; Arabidopsis sp

&lt;400&gt; 93

aattaaaatt tgagcgggtct aaaccattag accgtttaga gatccctcca acccaaaata  
 60  
 gtcgattttc acgtcttgaa catatattgg gccttaatct gtgtgggttag taaagacttt  
 120  
 tattgggtcaa agaaaaacaa ccatggccca acatgttgat acttttattt aattatacaa  
 180  
 gtaccctga attctctgaa atatatttga ttgaccaga tattaatttt aattatcatt  
 240



tccgtgtaaaa gtgaaggagt caccgtgact cgtcgtaatc tgaaaccaat ctgttcatat  
300  
gatgaagaag tttctctcgt tctcctccaa cgcgtagaaa attctgacgg cttaacgatg  
360  
tggcgaagat ctgttggtta tcgtttctct tcaagaatct ctgtttcttc ttcgttacca  
420  
aaccctagac tgattccttg gtcccgcgaa ttatgtgccg ttaatagctt ctcccagcct  
480  
ccggtctcga cggaatcaac tgctaagtta gggatcactg gtgttagatc tgatgccaat  
540  
cgagtttttg ccactgctac tgccgcccgt acagctacag ctaccaccgg tgagatttcg  
600  
tctagagttg cggctttggc tggattaggg catcactacg ctcgttgtta ttgggagctt  
660  
tctaaagcta aacttaggta tgtgtttact tttcttttct catgaaaaat ctgaaaattt  
720  
ccaattgttg gattcttaaa ttctcatttg ttttatgggt gtagtatgct tgtggttgca  
780  
acttctggaa ctgggtatat tctgggtacg ggaaatgctg caattagctt cccggggctt  
840  
tgttacacat gtgcaggaac catgatgatt gctgcatctg ctaattcctt gaatcaggtc  
900  
attgaaatgt tgagaagttc ataaatttcg aatccttggt gtgtttatgt agttgatctt  
960  
gcttgcttat gtttatgtag ttgaaaagtt taaaaatttc taatccttgg tagttgatct  
1020  
cgcttggttg ttttttcatt ttctagattt ttgagataag caatgattct aagatgaaaa  
1080  
gaacgatgct aaggccattg ccttcaggac gtattagtgt tccacacgct gttgcatggg  
1140  
ctactattgc tgggtgcttct ggtgcttggt tgttgccag caaggatgaat gtttgttttt  
1200  
ttatatgtga tttctttggt ttatgaatgg gtgattgaga gattatggat cttaaactttt  
1260  
gcttccacga caagggtatt gcagactaat atgttggtg ctggacttgc atctgccaat  
1320  
cttgactttt atgcgtttgt ttatactccg ttgaagcaac ttcacctat caatacatgg  
1380  
gttgccgctg ttgttggtgc tatccacccc ttgcttggtt aaatttttgt tccttttctt  
1440  
ctttatttta gcagattctg ttttggttga tactgctttt aattcaaaat gtagtcatgg  
1500  
ttcaccaatt ctatgcttat ctattttgtg tgttgtcagg tgggcggcag cgtctgggtca  
1560  
gatttcatac aattcgatga ttcttccagc tgctctttac ttttggcaga tacctcattt  
1620  
tatggccctt gcacatctct gccgcaatga ttatgcagct ggagggttaag accatatggt  
1680  
gtcatatgag attagaatgt ctccctccat gtagtggtga tcttgaacta gttcaatttc  
1740  
gtggaatgat cagagtgtcc tagatagtgt cacagcagtc gacattttag tggctagata  
1800  
atgagttctt tccgttagag ataaacattc gcgaacattg tttccagctt ccgcgaccca  
1860  
acttctgatt ttgtttcttg gtaccttggt ttcagttaca agatggtgct actctttgat  
1920  
ccgtcaggga agagaatagc agcagtggct ctaagggaact gcttttacct gatccctctc  
1980  
ggtttcatcg cctatgactg tgagtcttgt agattcatct tttttttgta gtttattgac  
2040  
tgcattgctg tatctgattt ttgctgttcc ttccaatttt tgtgacagaa ggggttaacct



2100  
caagttgggt ttgcctcgaa tcaacacttc tcacactagc aatcgctgca acagcatttt  
2160  
cattctaccg agaccggacc atgcataaag caaggaaaat gttccatgcc agtcttctct  
2220  
tccttctgt tttcatgtct ggtcttcttc tacaccgtgt ctctaattgat aatcagcaac  
2280  
aactcgtaga agaagccgga ttaacaaatt ctgtatctgg tgaagtcaaa actcagaggc  
2340  
gaaagaaacg tgtggctcaa cctccgggtgg cttatgcctc tgctgcaccg tttcctttcc  
2400  
tcccagctcc ttccttctac tctccatgat aacctttaag caagctattg aatttttggg  
2460  
aacagaaatt aaaaaaaaaa tctgaaaagt tcttaagttt aatctttggg taataatgaa  
2520  
gtggagaacg catacaagtt tatgtatttt ttctcatctc cacataattg tattttttct  
2580  
ctaagtatgt ttcaaattgat acaaaataca tactttatca attatctgat caaattgatg  
2640  
aatttttgag ctttgacgtg ttaggtctat ctaataaacg tagtaacgaa tttggttttg  
2700  
gaaatgaaat ccgataaccg atgatgggtg agagttaaac gattaaaccg ggttggttaa  
2760  
aggtctcgag tctcgacggc tgcggaaatc ggaaaatcac gattgaggac tttgagctgc  
2820  
cacgaagatg gcgatgaggt tgaaatcaat  
2850

<210> 94  
<211> 3660  
<212> DNA  
<213> Arabidopsis sp

<400> 94  
tatttgtatt tttattgtta aattttatga tttcaccgg tatatatcat cccatattaa  
60  
tattagattt attttttggg ctttatttgg gttttcgatt taaactgggc ccattctgct  
120  
tcaatgaaac cctaattgggt tttgtttggg ctttgattt aaaccgggcc cattctgctt  
180  
caatgaaggt cctttgtcca acaaaactaa catccgacac aactagtatt gccaagagga  
240  
tcgtgccaca tggcagttat tgaatcaaag gccgccaaaa ctgtaacgta gacattactt  
300  
atctccggta acggacaacc actcgtttcc cgaaacagca actcacagac tcacaccact  
360  
ccagtctccg gcttaactac caccagagac gattctctct tccgtcgggt ctatgacttc  
420  
gattctcaac actgtctcca ccatccactc ttccagagtt acctccgtcg atcgagtcgg  
480  
agtctctct cttcggaatt cggattccgt tgagttcact cgcgggcgtt ctggtttctc  
540  
gacgttgatc tacgaatcac ccggtagtta gcattctgtt ggatagattg atgaatgttt  
600  
tcttcgattt tttttttact gatcttggtg tggatctctc gtagggcgga gatttggtgt  
660  
gcgtgcggcg gagactgata ctgataaagg tatgattttt tagttgtttt tattttctct  
720  
ctcttcaaaa ttctcttttc aaacactgtg gcgtttgaat ttccgacggc agttaaatct  
780  
cagacacctg acaaggcacc agccgggtgg tcaagcatta accagcttct cggtatcaaa

840  
ggagcatctc aagaaactgt aattttgttc atctcctcag aatcttttaa attatcatat  
900  
ttgtggataa tgatgtgtta gtttaggaat tttcctacta aaggtaatct cttttgagga  
960  
caagtcttgt ttttagctta gaaatgatgt gaaaatgttg tttgtagct aaaaagagtt  
1020  
tgttgttata ttctgtattc agaataaatg gaagattcgt cttcagctta caaaaccagt  
1080  
cacttggcct ccactggttt ggggagtcgt ctgtgggtgct gctgcttcag gtaatcatat  
1140  
gaacctcttt tggatcatgc aatactgtac agaaagtttt ttcattttcc ttccaattgt  
1200  
ttcttctggc agggaaacttt cattggaccc cagaggatgt tgctaagtcg attctttgca  
1260  
tgatgatgtc tggtccttgt cttactggct atacacaggt ctggttttac acaacaaaaa  
1320  
gctgacttgt tcttattcta gtgcatttgc ttggtgctac aataacctag acttgtcgat  
1380  
ttccagacaa tcaacgactg gtatgataga gatatcgacg caattaatga gccatatcgt  
1440  
ccaattccat ctggagcaat atcagagcca gaggtaactg agacagaaca ttgtgagctt  
1500  
ttatctcttt tgtgattctg atttctcctt actccttaaa atgcaggtta ttacacaagt  
1560  
ctgggtgcta ttattgggag gtcttggtat tgctggaata ttagatgtgt gggtaagttg  
1620  
gcccttctga cattaactag tacagttaaa gggcacatca gatttgctaa aatcttccct  
1680  
tatcaggcag ggcataccac tccactgtc ttctatcttg ctttgggagg atcattgcta  
1740  
tcttatatat actctgctcc acctcttaag gtaagtttta ttctaactt ccactctcta  
1800  
gtgataagac actccatcca agttttggag ttttgaatat cgatatctga actgatctca  
1860  
ttgcagctaa aacaaaatgg atgggttgga aattttgcac ttggagcaag ctatattagt  
1920  
ttgccatggt aagatatctc gtgtatcaat aatatatggc gttgttctca tctcattgat  
1980  
ttgtttcttg ctcacttgac tgatagggtg gctggccaag cattgttttg cactcttacg  
2040  
ccagatgttg ttgttctaac actcttgtac agcatagctg gggtagctt ttggcaaacc  
2100  
ttttatgttg cttttttcgt tatctgttgt aatatgctct tgcttcatgt tgtaccttg  
2160  
tgataatgca gtttaggaata gccattgtta acgacttcaa aagtgttgaa ggagatagag  
2220  
cattaggact tcagtctctc ccagtagctt ttggcaccga aactgcaaaa tggatatgag  
2280  
ttggtgctat agacattact cagctttctg ttgccggtat gtactatcca ctgtttttgt  
2340  
gcagctgttg cttctatttc ttttccttga tcttatcaac tggatattca ccaatggtaa  
2400  
agcacaaatt aatgaagctg aatcaacaaa ggcaaacat aaaagtacat tctaataaaa  
2460  
tgagctaata aagaggaggc atctactttt atgtttcatt agtgtgattg atggattttc  
2520  
atttcatgct tctaaaacaa gtattttcaa cagtgtcatg aaataacaga acttatactt  
2580  
tcatttgtac ttttactagt ggatgagtta cacaatcatt gttatagaac caaatcaaag  
2640

gtagagatca tcattagtat atgtctatct tgggtgcagg atatctatta gcatctggga  
2700  
aaccttatta tgcgttggcg ttggttgctt tgatcattcc tcagattgtg ttccaggtaa  
2760  
agacgttaac agtctcacat tataattaat caaattcttg tcaactcgtct gattgctaca  
2820  
ctcgcttcta taaactgcag tttaaatact ttctcaagga ccctgtcaaa tacgacgtca  
2880  
agtaccagggt aagtcaactt agtacacatg tttgtgttct tttgaaatat ctttgagagg  
2940  
tctcttaatc agaagttgct tgaaacactc atcttgatta caggcaagcg cgcagccatt  
3000  
cttgggtgctc ggaatatttg taacggcatt agcatcgcaa cactgaaaaa ggcgtatttt  
3060  
gatgggggtt tgcgaaagc agagggtgtg acacatcaaa tgtgggcaag tgatggcatc  
3120  
aactagttta aaagattttg taaaatgtat gtaccgttat tactagaaac aactcctgtt  
3180  
gtatcaattt agcaaaacgg ctgagaaatt gtaattgatg ttaccgtatt tgcgctccat  
3240  
ttttgcattt cctgctcata tcgaggattg gggtttatgt tagttctgtc acttctctgc  
3300  
tttcagaatg tttttgtttt ctgtagtgga ttttaactat tttcatcact ttttgattg  
3360  
attctaaaca tgtatccaca taaaaacagt aatatacaaa aatgatactt cctcaaactt  
3420  
tttataatct aaatctaaca actagctagt aacccaacta acttcataca attaatgtga  
3480  
gaaactacaa agactagact atacatatgt tatttaacaa cttgaaactg tgttattact  
3540  
acctgatttt tttctattct acagccattt gatatgctgc aatcttaaca tatcaagtct  
3600  
cacgttggtg gacacaacat actatcacia gtaagacacg aagtaaaacc aaccggcaac  
3660